



Provided by the author(s) and University of Galway in accordance with publisher policies. Please cite the published version when available.

Title	Isolation and characterization of the bovine Stearoyl-CoAdesaturase promoter and analysis of polymorphisms in the promoter region in dairy cows
Author(s)	Keating, Aileen F.; Murphy, John J.; Smith, Terry J.; Cairns, Michael T.
Publication Date	2005-03
Publication Information	Keating AF, Stanton C, Murphy JJ, Smith TJ, Ross RP, Cairns MT. (2005) 'Isolation and characterization of the bovine Stearoyl-CoAdesaturase promoter and analysis of polymorphisms in the promoter region in dairy cows'. Mammalian Genome, 16 :184-193.
Publisher	Springer
Link to publisher's version	http://dx.doi.org/10.1007/s00335-004-2325-0
Item record	http://hdl.handle.net/10379/5815
DOI	http://dx.doi.org/10.1007/s00335-004-2325-0

Downloaded 2024-05-21T05:06:15Z

Some rights reserved. For more information, please see the item record link above.



Isolation and characterisation of the bovine *Stearoyl-CoA desaturase* promoter and analysis of polymorphisms in the promoter region in dairy cows.

Aileen F. Keating^{*}, Catherine Stanton[†], John J. Murphy[†], Terry J. Smith[°], R. Paul Ross[†] and Michael T. Cairns^{*}.

^{*}National Diagnostics Centre, National University of Ireland, Galway, Ireland.

[°]National Centre for Biomedical Engineering Science, National University of Ireland, Galway.

[†]Teagasc, Dairy Products Research Centre, Moorepark, Fermoy, Co. Cork, Ireland.

1

¹ Corresponding Author: Dr. Michael Cairns, National Diagnostics Centre, National University of Ireland, Galway. Email: Michael.cairns@nuigalway.ie.

The nucleotide sequence data reported in this paper have been submitted to GenBank and has been assigned the accession number AJ555480.

1 **ABSTRACT**

2
3 Conjugated linoleic acid (CLA) in milk arises through microbial biohydrogenation of
4 dietary polyunsaturated fatty acids (PUFA) in the rumen, and by the action of
5 mammary Stearoyl-CoA desaturase (Scd). A large variation (up to ten-fold) in the
6 concentration of this fatty acid in milk have been observed, even in cows receiving the
7 same diet. The reasons for this variation are not well understood. In this study, the
8 bovine core promoter region was isolated by a genome walking strategy from
9 genomic DNA GenomeWalker libraries and then cloned and characterised. This core
10 promoter sequence extended approximately 600bp upstream of the translation start
11 site. The presence of putative transcription factor binding sites conserved in bovine,
12 human, and mouse promoters were observed. Evidence that this promoter fragment
13 was functional *in vivo* was obtained from expression studies in a mammary cell line.
14 The promoter sequence of the *scd* gene was compared between cows selected for the
15 ability to produce high fatty acid methyl esters (FAME) (2.22-2.72) in their milk, with
16 the same promoter region of low FAME producing cows (0.81-1.12). However, such
17 comparisons of the sequences of the *scd* promoter region of cows producing high milk
18 CLA compared with low CLA revealed no polymorphisms in this promoter segment.
19 Furthermore, no sequence polymorphisms were observed between the *scd* promoter
20 region of Holstein Friesian, Montbeliarde, Normande, Norwegian Red, Charlois,
21 Limousin and Kerry breeds.

22
23 **(Key words:** Conjugated Linoleic acid, Stearoyl-CoA desaturase, Promoter,
24 Polymorphisms).

25
26 **Abbreviation key:** CLA = Conjugated Linoleic Acid, **scd** = Stearoyl-CoA
27 desaturase, **PCR** = polymerase chain reaction, **SNPs** = Single Nucleotide
28 Polymorphisms, **PUFA** = Polyunsaturated Fatty Acids, **FAME** = fatty acid methyl
29 esters.
30
31
32
33
34
35
36
37

38 **INTRODUCTION**

39

40 Conjugated Linoleic acid (CLA) is a collective term to describe one or more
41 positional and geometric isomers of linoleic acid (*cis*-9, *cis*-12-C_{18:2}). The *cis*-9,
42 *trans*-11-C_{18:2} isomer is the predominant one in the human diet as a result of microbial
43 biohydrogenation in the rumen. This isomer is produced in ruminants directly as an
44 intermediate during the microbial biohydrogenation of dietary linoleic acid, and
45 endogeneously from *trans*-vaccenic (*trans*-11-C_{18:1}) acid in mammary tissue by the
46 action of Stearoyl-CoA desaturase (Scd) (Griinari *et al.*, 2000). Animal-fat containing
47 foods including dairy products, beef and lamb are rich sources of CLA (Chin *et al.*,
48 1992; Fritsche and Steinhart, 1998; O’Shea *et al.*, 2000).

49

50 CLA has attracted much attention in recent years, due to its many potential health
51 benefits. Studies have shown that CLA exhibits anti-carcinogenic activity in animal
52 models (Belury *et al.*, 1995; Ha *et al.*, 1990; Ip *et al.*, 1996; Liew *et al.*, 1995) and in
53 *in vitro* studies using a range of human cancer cell lines including mammary (Miller
54 *et al.*, 2001; Park *et al.*, 2000; Visonneau *et al.*, 1997), prostate (Cesano *et al.*, 1998;
55 Palombo *et al.*, 2002) and colon, (Miller *et al.*, 2001; Palombo *et al.*, 2002). Other
56 properties of CLA include anti-atherogenic activity (Lee *et al.*, 1994; Nicolosi *et al.*,
57 1993), the ability to reduce the catabolic effects of immune stimulation (Cook *et al.*,
58 1993; Millar *et al.*, 1994), the ability to enhance growth promotion (Chin *et al.*, 1994)
59 and the ability to reduce body fat (Pariza *et al.*, 1996).

60

61 During biohydrogenation of linoleic acid in the rumen, CLA is produced. This CLA
62 is largely a transient intermediate and is involved as an intermediate in the
63 biohydrogenation pathway to produce stearic acid. A build up of *trans*-vaccenic acid

64 occurs (Harfoot and Hazelwood, 1988). *Trans*-vaccenic acid is also an intermediate
65 in the biohydrogenation of other PUFA (Griinari and Bauman, 1999). Several studies
66 have shown that substantial amounts of *trans*-vaccenic acid (60-300g/day) reach the
67 duodenum of lactating cows (Kalscheur *et al.*, 1997a, 1997b; Wonsil *et al.*, 1994).
68 Infusion studies using *trans*-vaccenic acid post-rationally resulted in elevated *cis*-9,
69 *trans*-11 CLA in milk-fat, indicating conversion in the mammary gland (Griinari and
70 Bauman, 1999). Several studies have found substantial Scd activity in both mammary
71 and adipose tissue of ruminant animals (Kinsella *et al.*, 1972; Martin *et al.*, 1999; St
72 John *et al.*, 1991, Ward *et al.*, 1998). Mammary Scd enzyme, which converts *trans*-
73 vaccenic acid to *cis*-9, *trans*-11 CLA is believed to be responsible for the formation of
74 the majority of CLA in milk (Griinari and Bauman, 1999).

75

76 Animal diet is a major factor affecting the milk-fat content of CLA, with dietary
77 supplements containing oils rich in PUFA, such as linoleic and linolenic acids, being
78 the most effective for CLA enrichment of milk. However, substantial variations in the
79 CLA content of milk-fat (ranging from 3 to 10-fold) of cows on the same dietary
80 treatment have been observed in a number of studies (Jiang *et al.*, 1996; Stanton *et al.*,
81 1997; Kelly *et al.*, 1998a; 1998b; Lawless *et al.*, 1998; 1999; Peterson *et al.*, 2002;
82 Solomon *et al.*, 2000; White *et al.*, 2001). The reasons for this variation are not well
83 understood. Perhaps ruminally derived CLA is a more important contributor to milk
84 CLA in grass-fed cows. The variation in the CLA content of milk-fat may also be due
85 to a variation in rumen conditions leading to differences in the availability of CLA (or
86 CLA precursors) that escape from the rumen. It might also be caused by differences in
87 mammary Scd activity associated with either regulation of *scd* gene expression,
88 differences in structure of the enzyme due to gene polymorphisms, or differences in

89 downstream factors that would affect interaction between enzyme and substrate (e.g.
90 phosphorylation) (Peterson *et al.*, 2002).

91
92 The promoter region of the human (Bene *et al.*, 2001; Zhang *et al.*, 2001), chicken
93 (Lefevre *et al.*, 2001), and mouse (Ntambi *et al.*, 1988; Kaestner *et al.*, 1989; Mihara,
94 1990) *scd* genes have been isolated, cloned and characterized. It has been shown in
95 these studies that there is a conserved PUFA response region in all three, and that this
96 includes critical binding sites for Sterol Response Element Binding Protein (SREBP)
97 and Nuclear Factor-Y (NF-Y) transcription factors. Sequence comparison of the
98 human and mouse promoters indicated a second region of high homology including
99 the 5'UTR and basal/proximal promoter. It has been suggested that there are two
100 different transcription start sites in the human promoter, and that these may be
101 dependent on tissue-specific factors (Zhang *et al.*, 2001).

102
103 In this study, the role that polymorphisms in the core promoter region of the bovine
104 *scd* gene might play in influencing regulation of the *scd* gene was investigated. The
105 bovine core promoter was therefore isolated from genomic DNA by a genome
106 walking approach using primers designed to a genomic database sequence for the
107 bovine *Stearoyl-CoA desaturase* gene (AF481915) that contained 162 bases of
108 5'UTR. Analysis of this promoter sequence identified a number of conserved
109 potential transcription factor binding sites based on comparison with the human and
110 mouse *scd* promoter regions and interrogation of the TRANSFAC 4.0 database of
111 transcription factor sequences (Wingender *et al.*, 2000) with the MatInspector V2.2
112 program (Quandt *et al.*, 1995). Importantly, and for the first time for bovine *scd*, this
113 core promoter was shown to drive transcription of the reporter gene luciferase in an *in*

114 *vitro* mammalian culture system, confirming its presumed function. A number of
115 animals were screened for possible polymorphisms in this promoter region. These
116 included cows producing high milk-fat CLA as a % of fatty acid methyl ester (%
117 FAME) content (2.22-2.72), compared with cows yielding a low milk-fat CLA as a %
118 of FAME content (0.81-1.12). Additionally, sequence comparisons of the *scd*
119 promoter region from a number of different breeds were made.
120 A longer bovine *scd* promoter sequence (Acc No. AY241932) is now included in the
121 Genbank database but this was not available at the time of this study: the two
122 promoter sequences were submitted within a very short time of each other.

123 **MATERIALS AND METHODS**

124

125 **Construction of GenomeWalker (GW) libraries**

126 To obtain the 5' flanking sequence of the *scd* gene, GenomeWalker (GW) libraries
127 were constructed from bovine total genomic DNA using a Universal GenomeWalker
128 kit (Clontech, UK) according to the manufacturer's instructions. High quality
129 genomic DNA was first extracted from cultured lymphoblast cells from a Holstein-
130 Friesian animal. Briefly, a cell suspension (15 ml) was centrifuged at 1200 g for 5min
131 and the pellets mixed and incubated overnight at 37°C in 900 µl 0.2M EDTA 0.5%
132 sodium-*n*-lauroyl sarcosine and 25µl of proteinase K (20mg/ml). 10µl of RNase
133 (2mg/ml) was added to each tube and incubated at 37°C for 1.5 hr. The mix was split
134 in two, 200 µl phenol added, mixed and incubated for 30 min at 37°C in a rotary
135 mixer. Chloroform (200 µl) was added, tubes were shaken vigorously and incubated
136 at 37°C for 1 hr in a rotary mixer. Tubes were centrifuged at 16000 g for 15 min to
137 form two layers. The upper layer was transferred to a fresh tube and two volumes of
138 ice-cold 100% ethanol added. Tubes were inverted abruptly four times, and at this
139 point a DNA precipitate was obtained. The supernatant was decanted, 100 µl 70%
140 ethanol added and incubated overnight at 21°C in a rotary mixer. Ethanol was
141 decanted off and the pellet was allowed to air-dry. The DNA pellets were
142 resuspended in 50 µl 10mM Tris-HCl, pH8.0 containing 1mM EDTA (TE) buffer and
143 aliquots were checked for integrity on a 0.7% w/v agarose gel containing ethidium
144 bromide (EtBr) (2µg/ml) gel.

145

146 The genomic DNA was digested with four restriction enzymes, *DraI*, *StuI*, *EcoRV*,
147 and *PvuII*, to create four pools of DNA fragments. Adaptors were ligated onto both

148 ends of these fragments to create four GW libraries and these libraries were
149 designated DL1 (*DraI*), DL2 (*StuI*), DL3 (*EcoRV*), and DL4 (*PvuII*).

150

151 **Isolation and characterisation of the *scd* promoter from GW libraries**

152 To specifically isolate the *scd* promoter, genome walking primers pAKGW1 (5'-
153 GCTCTCAGACACTGGGATCACTTTCTCGGG-3') and pAKGW2 (5'-
154 AACTGAGTGTAGACTAGTTCCTGAGCCTGC-3') were designed using Vector
155 NTI software (InforMax, Inc.) to a genomic database sequence for the bovine *scd*
156 gene (AF481915) and synthesized by MWG Biotech (Ger). The primary PCR was
157 carried out using 1µl of the GenomeWalker libraries with the gene-specific primer,
158 pAKGW1, and adaptor primer, AP1 (from kit). The primary PCR products were
159 diluted 1:50 and used as template for nested PCR with a second gene-specific primer,
160 pAKGW2, and a second adaptor primer AP2 (again from kit). The reaction final
161 volume was 50 µl containing *Taq* DNA polymerase buffer (Invitrogen), 1.5mM
162 MgCl₂, 200µM dNTPs (Promega), 0.3µM each primer, and 1U *Taq* polymerase
163 (Invitrogen). Immediately prior to cycling 2.5 µl of DMSO was added to the mix.
164 The reaction was amplified for 35 cycles for both the first and second PCR. Cycling
165 was performed in a DNA Engine thermal cycler (MJ Research) and conditions were
166 95°C for 2min, followed by 35 cycles of 95°C for 30s, 60°C for 30s and 72°C for 3
167 min. This was followed by a 72°C final extension step for 7 min.

168

169 PCR products were directly cloned into the pCR2.1 vector (Invitrogen) and
170 transformed using One Shot Top-10 chemically competent *E. coli* cells according to
171 the manufacturer's instructions (Invitrogen). Colony PCR was performed to identify
172 clones that potentially carried the *scd* promoter. Small-scale preparations of plasmid

173 DNA were made from these colonies using the procedure outlined by the
174 manufacturer (Sigma). Sequencing of three plasmids (DL1.2, DL1.4 and DL1.8) was
175 performed by MWG Biotech (Ger). The resulting sequences were analysed using
176 Vector NTI software (InforMax, Inc). Potential transcription factor binding sites were
177 identified using the MatInspector V2.2 program (Quandt *et al.*, 1995) by interrogation
178 of the TRANSFAC 4.0 database of transcription factor sequences (Wingender *et al.*,
179 2000).

180

181 **Luciferase reporter vector construction**

182 pCR2.1-based plasmids containing the putative *scd* promoter were restriction digested
183 at 37°C to release the fragment and create compatible ends for sub-cloning into the
184 pGL3-Basic (Promega) promoter-less expression vector. The pGL3-Basic vector was
185 digested in the same manner. Ligation reactions were transformed into TAM
186 Ultracomp chemically competent *E. coli* cells (Active Motif Europe), using
187 instructions recommended by the manufacturer.

188

189 **Cell culture**

190 Chinese Hamster Ovary K1 (CHO-K1) cells obtained from ATCC (CCL-61) were
191 cultured in Dulbecco's Modified Eagles Medium F12 (Biowhittaker, UK) containing
192 10% (v/v) foetal bovine serum (Invitrogen). Human mammary MCF-7 cells (ATCC –
193 HTB22) were cultured in Eagles Minimum Essential Media (Invitrogen) containing
194 10% (v/v) foetal bovine serum (Invitrogen) and 1% (v/v) non-essential amino acids
195 (Invitrogen). Cells were routinely passaged every 3-4 days by washing with
196 phosphate buffered saline (PBS) and treating with 3ml EDTA-trypsin (Sigma) to
197 remove adherent cells. Cells were maintained in a humidified incubator at 37°C and

198 5% CO₂. Cells were grown to 80% confluency and both CHO-K1 and MCF-7 cells
199 were transiently co-transfected with luciferase reporter vectors using Fugene 6
200 transfection reagent (Roche Diagnostics). Six-well transfection plates were seeded at
201 a concentration of 3 x 10⁵ cells per well and incubated overnight at 37°C and 5% CO₂.
202 pGL3-Scd experimental constructs (1µg) were co-transfected with 25ng of pRL-SV40
203 plasmid (Promega) to control for transfection efficiency. Forty-eight hours after
204 transfection, media was removed from the wells, 300µl Passive Lysis Buffer
205 (Promega) was added followed by incubation at room temperature for 10min.

206

207 **Luciferase assay**

208 The Dual Luciferase Assay Kit (Promega) was used to measure both Renilla and
209 Firefly luciferase expression by the reporter vectors on a Tecan Spectrafluor Plus
210 luminometer using the Magellan software (Tecan). Mean Firefly luciferase activity
211 values were corrected for variations in transfection efficiency using the corresponding
212 mean Renilla luciferase figures. The corrected values were expressed as a percentage
213 of the positive control value (pGL3-Control). Transfection values were a result of
214 three independent transfections, with n=6, for both cell types.

215

216 **Analysis of *scd* promoter polymorphism incidence**

217 A dairy herd (n=75) had been on a ryegrass diet for a period of six months and their
218 milk was analysed for fatty acid methyl esters content (% FAME) using the gas liquid
219 chromatography (GLC) method described previously (Stanton *et al.*, 1997). FAME
220 content was measured in milk samples from 59 animals taken at the evening milking
221 on two occasions during the grazing season, in July and September. Nine of these 75
222 cows with consistently low and high milk CLA as a % of FAME values over both

223 sampling times were selected for this study. The CLA isomer measured was the *cis*-9,
224 *trans*-11 CLA isomer. High CLA animals produced greater than 2.0% FAME (2.22-
225 2.72) in their milk, compared with low milk CLA producers of ~ 1.0% FAME (0.81-
226 1.12). The CLA as a % of FAME ranged from 0.81 to 2.72 as quoted in Table I and
227 are the mean values of both sampling times.

228

229 Genomic DNA, for PCR amplification of the *scd* promoter fragment, was extracted
230 from whole blood (200µl) collected in heparinised tubes from the coccygeal vein from
231 these nine Holstein Friesian cows and in addition ten cows of different bovine breeds
232 using a Gentra capture column™ (Gentra). Amplification of the bovine promoter was
233 performed using primers D9Dfor (5'-TGATGGGGTAGTGAGGAGC-3') and
234 D9Drev (5'-G TTCCTGAGCCTGCTTTTGC-3') with 1µl of genomic DNA as
235 template (~200ng), in a final volume of 50µl, containing *Taq* DNA polymerase buffer
236 (Invitrogen), 1.5mM MgCl₂, 200µM dNTPs (Promega), 0.3µM each primer, and 1U
237 *Taq* polymerase (Invitrogen). Cycling was performed in a DNA Engine thermal
238 cycler (MJ Research) and conditions were 95°C for 1min, followed by 35 cycles of
239 93°C for 1min, 58.5°C for 1min, 72°C for 2min and a final extension step of 72°C for
240 10min. PCR products were purified using a PCR purification kit (Qiagen). DNA was
241 eluted in 30µl of PCR grade water (Sigma).

242

243 **RESULTS and DISCUSSION**

244

245 **Isolation of the bovine *scd* promoter**

246 Isolation of the bovine *scd* promoter was achieved using a genome walking strategy,
247 which involved digestion of bovine genomic DNA with four restriction enzymes,
248 *Dra*I, *Stu*I, *Eco*RV, and *Pvu*II, to create pools of short DNA fragments of varying
249 lengths. Adaptors were ligated onto the ends of these pools of DNA and PCR
250 employed to “walk” along the genome. Primers for PCR were designed to a bovine
251 genomic sequence for the *scd* gene (AF481915) that extends approximately 160bp
252 upstream of the translation initiation codon. PCR products were obtained for libraries
253 DL1 (*Dra*I), DL2 (*Stu*I) and DL3 (*Eco*RV), but not for the DL4 (*Pvu*II) library.
254 Library DL1 yielded a mixture of fragments but as these were also the longest, these
255 fragments were purified as a mixed pool and cloned into the pCR2.1 vector. Colony
256 PCR was performed and three plasmids (DL1.2, DL1.4 and DL1.8) were sequenced.
257 BlastN analysis of the three sequences indicated that they were all fragments of the
258 bovine *scd* gene and the DL1.4 sequence extended upstream of the bovine genomic
259 database sequence, AF481915.

260

261 **Comparison of bovine, human and mouse *scd* promoters**

262 An alignment of the DL1.4 sequence was performed with a human promoter sequence
263 (AF320307) (Zhang *et al.*, 2001) and a mouse promoter sequence (M21280) (Figure
264 I). The bovine sequence displayed 67% identity with the human *scd* sequence and
265 59% homology with the murine promoter sequence *scd1*. The bovine sequence has
266 been submitted to the Genbank database (Acc. No. AJ555480). Transcription start
267 sites have been identified in both the mouse (Ntambi *et al.*, 1988) and human (Zhang

268 *et al.*, 2001) genes at 28bp and 37bp respectively from the proximal TATA box in the
269 bovine sequence. The bovine promoter sequence presented in Figure I is numbered
270 relative to the human transcription start site, and extends to -407.

271

272 Potential transcription factor binding sites were compared with those predicted for the
273 human promoter (Zhang *et al.*, 2001). Putative transcription-factor binding sites
274 identified using the MatInspector program (Quandt *et al.*, 1995) included Nuclear
275 factor Y, Octamer binding factor 1, and muscle-specific Mt-binding sites.
276 Comparison of the mouse, human and bovine sequences identified a number of
277 conserved sequences in two particular regions of the promoters. The first area of
278 conservation occurs between -37 to -119, where there are two conserved TATA
279 sequences (5'-TTTAAAT-3' and 5'-TAAAA-3'), a fat-specific element (FSE) (5'-
280 CTGAGGAAA-3') (-77 to -86), and binding sites for the transcription factors AP-1 (-
281 64 to - 68), NF-1 (-102 to -109) and HNF4 (-116 to -119). Analysis of the human
282 *SCD* promoter has shown that for liver and hair follicles, there was a major
283 transcription initiation site 35 nucleotides downstream from the proximal TATA box
284 (Zhang *et al.*, 2001). It was also shown that another transcription initiation site was
285 present 37 nucleotides downstream from the distal TATA box (Bene *et al.*, 2001), and
286 it was suggested that the human *SCD* gene has different start sites that depend on
287 different tissue-specific factors (Zhang *et al.*, 2001). The TATA sequence,
288 TTTAAAT, is somewhat unusual: this sequence, where the A in the second position
289 is replaced by a C, T, or G, has been shown to reduce the efficiency of a promoter in
290 *in vitro* studies (Conchino *et al.*, 1983).

291

292 The second conserved region occurs at -313 to -390 and has been designated the
293 PUFA response region. Putative binding sites for transcription factors NF-Y (-313 to
294 -317), NF-1 (-333 to -336), NF-Y/NF-1 (-351 to -361), SREBP (-366 to -376), and
295 SP-1 (-386 to -390) are conserved between the three promoter sequences. The PUFA
296 response region present in the bovine sequence is an element previously shown to
297 mediate the down-regulation of mouse *scd* expression in response to PUFA (Waters *et*
298 *al.*, 1997). The Sterol Response Element binding protein (SREBP) has been shown to
299 activate *scd* genes in the mouse (Tabor *et al.*, 1999; Shimomura *et al.*, 1998), and it
300 has also been shown that PUFA negatively regulate SREBP mRNA and protein
301 activation (Xu *et al.*, 1999; Yahagi *et al.*, 1999). This bovine promoter fragment also
302 contains the CCAAT box at -356 to -360 previously shown to be critical for
303 transcriptional activation in the human promoter (Zhang *et al.*, 2001). This CCAAT
304 element is also required for full activation of mouse *scd1* and *scd2* promoters and
305 binds the NF-Y transcription factor (Tabor *et al.*, 1999).

306

307 **Transcription studies in CHO-K1 and MCF-7 cells**

308 To determine whether this 407bp putative promoter sequence was sufficient to direct
309 transcription, luciferase reporter gene constructs containing the putative promoter
310 cloned into the promoter site of the promoter-less expression vector pGL3-Basic were
311 constructed. A plasmid (pRL-SV40) providing constitutive expression of Renilla
312 luciferase was co-transfected to serve as transfection efficiency control. Two cell
313 types were transfected, a human mammary cell line, MCF-7, and a Chinese hamster
314 ovary cell line, CHO-K1. The CHO-K1 cell line is most commonly used to express
315 mammalian genes *in vitro* and should contain the general transcription factors
316 necessary for activating transcription of most genes. The human mammary cell line

317 contains more tissue-specific transcription factors, and is a more suitable system for
318 studying regulation of a gene expressed in the mammary gland.

319

320 The putative promoter showed promoter activity in both CHO-K1 and MCF-7 cell
321 lines (Figure II). Promoter activity was greatest in the mammary MCF-7 cell line
322 being 6-fold higher in MCF-7 cells than in CHO-K1 cells. These results indicate that
323 the 407bp region upstream of the proposed transcription start site is sufficient to direct
324 transcription.

325

326 **Promoter polymorphism screen**

327 Animals receiving an identical dietary treatment of ryegrass for a 6-month period
328 were seen to have milk CLA as a % of FAME varying from 0.81 to 2.72. The sample
329 size selected was initially 59 Holstein Friesian cows, from which milk was analysed
330 for CLA content and subsequently 9 cows were selected and divided into two groups
331 of high and low CLA producers. There were no differences in the milk yields, feed
332 intake and milk fat content of the high v's low CLA animals. However, we did
333 observe that blood glucose and non-essential fatty acids (NEFA) differed significantly
334 between high and low CLA cows, with glucose being higher in the blood of the low
335 CLA group whereas NEFA was lower in the low CLA group compared to the high
336 CLA group. To investigate whether polymorphisms in the bovine *scd* promoter could
337 explain the considerably large inter-cow variation in milk fat CLA content, the
338 nucleotide sequences of the promoter region of nine cows (Table I) were analysed to
339 identify sequence polymorphisms which may be related to elevation of milk fat CLA.

340

341 In an attempt to also include an increased genetically diverse element, the promoter
342 region of the *scd* gene was amplified and sequenced from ten animals of different
343 breeds (high genetic merit Holstein Friesian, low genetic merit Holstein Friesian,
344 Montbeliarde, Normande, Norwegian Red, Irish Friesian, Dutch Freisian, Charlois,
345 Limousin, Kerry). No information regarding the milk CLA status of these animals
346 was available.

347

348 The *scd* promoter region sequences were aligned for these nineteen animals (nine
349 animals showing high and low milk CLA as a % of FAME and ten animals of varying
350 bovine breeds) and compared to search for any polymorphisms in the area (Data not
351 shown). This alignment showed the total absence of any polymorphic sites between
352 the bovine *scd* promoters of 19 (nine high and low milk CLA and ten animals of
353 different breeds). This high conservation may be significant in indicating that the
354 regulation of this gene is under extremely rigid control by transcription factors, and a
355 high conservation is necessary for full activation.

356

357 **CONCLUSION**

358

359 In conclusion we have isolated and partially characterised the bovine *scd* promoter.
360 This promoter fragment showed 59% and 67% similarity to both the mouse and
361 human sequences, respectively. Differences in the CLA content of milk from these
362 animals were not a consequence of polymorphisms within the core promoter of this
363 gene. The *scd* promoter sequence exhibits remarkable sequence conservation not just
364 across high and low CLA yielding Holstein Friesians but also across ten different
365 breeds. In contrast considerable differences were seen between human and mouse
366 sequences which included variation in non-binding regions and binding regions for

367 AP-1, SP-1, and NF-KB transcription factors. Consequently the observed variation in
368 the levels of milk CLA produced in the Holstein Friesian animals may be explained
369 by differences in ruminant synthesis of CLA or CLA precursors, polymorphisms in
370 the coding sequences of the bovine *scd* gene, or differences in the regulatory proteins
371 themselves, an area that requires further investigation.

372

373 **ACKNOWLEDGEMENTS:**

374 This work was supported by a Teagasc Walsh Fellowship. The human mammary cell
375 line, MCF-7, was a gift from Dr. C Curran, University College Hospital, Galway.

376 **TABLES and FIGURES**

377 Table I: High and low CLA animals.

Animal	Breed	% FAME
330	Holstein Friesian	0.81 +/- 0.49
385	Holstein Friesian	0.88 +/- 0.49
387	Holstein Friesian	1.04 +/- 0.49
44	Holstein Friesian	1.12 +/- 0.49
728	Holstein Friesian	2.22 +/- 0.49
67	Holstein Friesian	2.27 +/- 0.49
249	Holstein Friesian	2.32 +/- 0.49
376	Holstein Friesian	2.34 +/- 0.49
225	Holstein Friesian	2.72 +/- 0.49

378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403


404 **Figure I: Alignment of Human and Mouse Stearoyl-CoA desaturase promoters**
 405 **with DL1.4 (Bovine) sequence.**
 406

```

407
408
409 Human          -407                -390
410 Mouse          -GGGGGAGCGAGGAGCTGGCGGCAGAGGGAACAGCAGATTGCGCCGAGCCAATGGCAACG
411 Bovine          GGAGAGACGGAGAAGCTAGAGGCAGAGGGAACAGCAGATTGCGCCTAGCCAATGGAAAAG
412                -GGGGTAGTGAGGAGCTCGCGGCAGAGGGAACATCAGATTGCGCCGAGCCAATGGCAACG
413                * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *
414                * * * * * * * * * * * * * * * * * * * * * * * * * * * *
415                Sp-1          RFX1          SREBP          NF-Y/NF-1/RFX1
416
417 Human          -313
418 Mouse          GCAGGACGAGGTGGCACCAAATTCCCTTCGGCCAATGACGAGCCGGAGTTTACAGAAGCC
419 Bovine          GCAGGACAAGTGGCACCAAATTCTTTGGCCAATGACAAGACGGGCTTCACAGGAGGC
420                GCAGGACGAGGTGGCACCAAATTCCCTTCGGCCAATGACGCGCCAGAGTCTACAGAAGCC
421                *****
422                NF-1          NF-Y/AP-1
423
424 Human          TCATTAGCATTTCCCAGAGGCAGGGG-CAGGGGCAGAGG--CCGGGTGGTGTG-----
425 Mouse          ACATTAGCATTTATCCCAGGCAGGGGTTGGAGCAGCGCCCTGTTGTATGCCCTTCAGC
426 Bovine          -CATTAGCATTTCCCAGGGGCAGGGG-CAGAGGCAGGGG---CTGGCGCGGCC----AA
427                ***** * * * * * * * * * * * * * * * * * * * * * *
428
429 Human          GTGTCCGTGTCGGCAGCATC--CCCGCGCCCTGCTGCGGT-----CGCCCGAGCC
430 Mouse          ATCCCAGCGCCTCCAAGGTCTACTCTGGAATCTACTTGGCTTTCTTTCCCGTTCCTGGTC
431 Bovine          GCCGCGGTGTGTGCAGCA---TCCAGTCTTGCTTCTTC-----GGCCCCAG-C
432                * * * * * * * * * * * * * * * * * * * * * * * * * * * *
433
434 Human          TCGGCCTCTGTCTCCTCC-----CCCTC
435 Mouse          CCGCCCTCTCTCTCCTCCCTCCCTCCCTCCCTCCCTCCCTCCCTCCCTCCCTCCCTC
436 Bovine          ACGCCTCGGCGCTCTGTC-----TCCTC
437                * * * * * * * * * * * * * * * * * * * * * * * * * * * *
438
439 Human          -119
440 Mouse          CCGCCCTTACCTCCACGCGGGACCGCCGCGCCAGTCAACTCCTCGCACTTTGCCCTTGC
441 Bovine          CCTCCCTCACCTCCACGCCTGGCTTCCCTGGCTAGCTATCTCTGCGCTCTTTACCCTTTG
442                CC-CTCTCCCGCCCATGCGGATCTCCACGGTGAACCAACTCTGCGCACTTTGCCCTTGC
443                * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *
444                HNF4
445
446 Human          TTGGCAGCGGATAAAAAGGGGGCTGAGGAAATACCGACACGGTCA-CCC GTT GCCAGCTC
447 Mouse          CTGGCAGCCGATAAAAAGGGGGCTGAGGAAATACTGAACACGGTCAATCCCATCGCCTGCTC
448 Bovine          TTGGCAACGAAATAAAAAGAGGTCTGAGGAAATACGGGACACAGTCA-CCCCGTGCCAGCGC
449                ***** * * * * * * * * * * * * * * * * * * * * * * * * * * * *
450                NF-1          TATA          FSE          AP-1
451
452 Human          -37
453 Mouse          TAGCCTTTAAATTCCTGGCTCGGG-GACCTCCACGCACCGCGGCTAGCGCCGACAACCAG
454 Bovine          TACCCTTTAAATTCCTGGCTCGGG-GACCTCCACGCACCGCGGCTAGCGCCGACAACCAG
455                TAGCCTTTAAATTCCTGGCTCGGG-GACCTCCACGCACCGCGGCTAGCGCCGACAACCAG
456                * * * * * * * * * * * * * * * * * * * * * * * * * * * *
457                TATA
458
459 Human          +1
460 Mouse          CTAGCGTGCAAGG---CGCCGCGGCTC---AGCGC-GTACCGGCGGGCTTCGAAACCGCA
461 Bovine          CCCGAGAGTCAGG---AGGGCAGGTTTCCAAGCGAGTTCCGCCACTCGCCTACCCAAC
462                CCAGCGCGCACCGTGCAGCGGAAGTCCCAGCGCAGCGCTGCGGATC-CCCACGCAAA-
463                * * * * * * * * * * * * * * * * * * * * * * * * * * * *
464
465 Human          GTCCTCCGGCGACCCCGAACTCCGCTCCGGAGCCTCAGCCCCCTGGAAAGTG
466 Mouse          GGGCTCCGGAA---CCGAAGTCCACGCTCGA-TCTCAGCACTG-GGAAAGTG
467 Bovine          -----


```

466 **Figure II: Transient transfection of mammalian cell lines with promoter**
467 **constructs.**



484 (a)


CHO-K1 cells



485

486 (b)

MCF-7 cells



487

488 **FIGURE LEGENDS**

489

490 Figure I: Sequence alignment of the DL1.4 sequence with database entries for the
491 human (AF320307) and murine (M21280), *scd1* promoters. Numbering is indicated
492 above sequence. Transcription factor binding sites are underlined and bold and these
493 sites are based on TRANSFAC 4.0 database search results and comparison of human
494 and mouse *scd* gene promoters. The critical CCAAT box from human studies is
495 overlined. Numbering is relative to the human transcription start site and is indicated
496 by +1. The transcription start site in the mouse promoter is indicated by the symbol ●.
497 Abbreviations: RFX-1=X-box-binding regulatory factor 1, SREBP=Sterol response
498 element binding protein, NF-Nuclear factor.

499

500 Figure II: Transient transfection of mammalian cells with the putative bovine *scd*
501 promoter sequence. (a) Luciferase assay following transient co-transfection of CHO-
502 K1 cells with promoter construct. (b) Luciferase assay following transient co-
503 transfection of MCF-7 cells with promoter construct. Luciferase activities are given
504 as a ratio of Firefly to Renilla (pRL-SV40) values

505 **REFERENCES**

506 Belury, M.A. 1995. Conjugated dienoic linoleate: a polyunsaturated fatty acid with
507 unique chemoprotective properties. *Nutr. Rev.* **53**: 83-89.

508

509 Bene, H., Lasky, D. and Ntambi, J.M. 2001. Cloning and characterisation of the
510 human Stearoyl-CoA desaturase gene promoter: Transcriptional activation by sterol
511 regulatory element binding protein and repression by polyunsaturated fatty acids and
512 cholesterol. *Biochem. Biophys. Res. Comm.* **284**: 1194-1198.

513

514 Cesano, A., Vissonneau, S., Scimeca, J.A., Kritchevsky, D. and Santoli, D. (1998).
515 "Opposite effects of linoleic acid and conjugated linoleic acid on human prostate
516 cancer in SCID mice." *Anticancer Res.* **18**: 1429-1434.

517

518 Chin, S.F., Liu, W., Storkson, J.M., Ha, Y.L. and Pariza, M.W. 1992. Dietary
519 sources of conjugated dienoic isomers of linoleic acid, a newly recognised class of
520 anticarcinogens. *J. Food. Comp. Anal.* **5**: 185-197.

521

522 Chin, S. F., Storkson, J.M., Albright, K.J., Cook, M.E. and Pariza, M.W. (1994).
523 "Conjugated linoleic acid is a growth-factor for rats as shown by enhanced weight
524 gain and improved feed efficiency." *J. Nutr.* **124**: 2344-2349.

525

526 Conchino, M., Goldman, R.A., Caruthers, M.H. and Weinmann, R. (1983). "Point
527 mutations of the adenovirus major late promoter with different transcriptional
528 efficiencies in vitro." *J. Biol.Chem.* **258**: 8493-8496.

529 Cook, M.E., Miller, C.C., Park, Y. and Pariza, M.W. 1993. Immune modulation by
530 altered nutrient metabolism: nutritional control of immune-induced growth
531 depression. *Poult. Sci.* **72**: 1301-1305.

532

533 Fritsche, J. and Steinhart, H. 1998. Amounts of conjugated linoleic acid (CLA) in
534 German foods and evaluation of dairy intake. *Z. Lebensm. Unters. Forsch. A.* **206**:
535 77-82.

536

537 Griinari, J.M. and Bauman, D.E. 1999. Biosynthesis of conjugated linoleic acid and
538 its incorporation into meat and milk in ruminants. *Advances in conjugated linoleic*
539 *acids research.* M. P. Yurawecz, Mossaba, M.M., Kramer, J.K.G., Pariza, M.W.,
540 Nelson, G.J. IL, AOCS Press. **1**: 180-200.

541

542 Griinari, J. M., Corl, B.A., Lacy, S.H., Chouinard, P.Y., Nurmela, K.V. and Bauman,
543 D.E. 2000. Conjugated linoleic acid is synthesised endogenously in lactating cows
544 by Delta-9 desaturase. *J. Nutr.* **130**: 2285-2291.

545

546 Ha, Y.C., Storkson, J.M. and Pariza, M.W. 1990. Inhibition of benzo (a) pyrene-
547 induced mouse forestomach neoplasia by conjugated dienoic derivatives of linoleic
548 acid. *Cancer Res.* **50**: 1097-1101.

549

550 Harfoot, C. G., and Hazlewood, G.P. 1988. Lipid metabolism in the rumen. The
551 rumen microbial ecosystem. P. N. Hobson. London, Elsevier Applied Science
552 Publishers 285-322.

553

554 Ip, C., Briggs, S.P., Haegele, A.D., Thompson, H.J., Storkson, J. and Scimeca, J.A.
555 (1996). "The efficacy of conjugated linoleic acid in mammary cancer prevention is
556 independent of the level or type of fat in the diet." Carcinogenesis. 17: 1045-1050.
557

558 Jiang, J., Bjoerck, L., Fonden, R. and Emanuelson, M. 1996. Occurance of
559 conjugated *cis*-9, *trans*-11-octadecadienoic acid in bovine milk: Effects of feed and
560 dietary regimen. J. Dairy Sci. **79**: 438-445.
561

562 Kaestner, K. H., Ntambi, J.M., Kelly Jr, T.J. and Lane, M.D. 1989. Differentiation-
563 induced gene expression in 3T3-L1 preadipocytes. A second differentially expressed
564 gene encoding Stearoyl-CoA desaturase. J. Biol. Chem. **264**: 14755-14761.
565

566 Kalscheur, K.F., Teter, B.B., Piperova, L.S. and Erdman, R.A. 1997a. Effect of
567 dietary forage concentration and buffer addition on duodenal flow of *trans*-C_{18:1} fatty
568 acids and milk fat production in dairy cows. J. Dairy Sci. **80**: 2104-2114.
569

570 Kalscheur, K.F., Teter, B.B., Piperova, L.S. and Erdman, R.A. 1997b. Effect of fat
571 source on duodenal flow of *trans*-C_{18:1} fatty acids and milk fat production in dairy
572 cows. J. Dairy Sci. **80**: 2115-2126.
573

574 Kinsella, J.E. 1972. Stearoyl-CoA as a precursor of oleic acid and glycerolipids in
575 mammary microsomes from lactating bovine: possible regulatory step in milk
576 triglycerides synthesis. Lipids. **7**: 349-355.
577

578 Kelly, M.L., Berry, J.R., Dwyer, D.A., Griinari, J.M., Chouinard, P.Y., Van
579 Amburgh, M.E. and D.E. Bauman. 1998a. Dietary fatty acid sources affect
580 conjugated linoleic acid concentrations in milk from lactating dairy cows. J. Nutr.
581 **128**: 881-885.

582

583 Kelly, M.L., Kolver, E.S., Bauman, D.E., Van Amburgh, M.E. and Muller, L.D.
584 1998b. Effect of intake of pasture on concentrations of conjugated linoleic acid in
585 milk of lactating cows. J. Dairy. Sci. **81**: 1630-1636.

586

587 Lawless, F., Murphy, J.J., Harrington, D., Devery, R. and Stanton, C. 1998.
588 Elevation of conjugated *cis*-9, *trans*-11-Octadecadienoic acids in bovine milk because
589 of dietary supplementation. J. Dairy Sci. **81**: 3259-3267.

590

591 Lawless, F., Murphy, J.J., Darrington, D, Devery, R, and Stanton, C. (1999).
592 “Influence of breed on bovine milk *cis*-9, *trans*-11 conjugated linoleic acid content.”
593 Livestock. Prod. Sci. 62: 43-49.

594

595 Lee, K.N., Kritchevsky, D. and Pariza, M.W. 1994. Conjugated linoleic acid and
596 atherosclerosis in rabbits. Atherosclerosis. **108**: 19-25.

597

598 Lefevre, P., Tripon, E., Plumelet, C., Douaire, M. and Diot, C. 2001. Effects of
599 Polyunsaturated Fatty Acids and Clofibrate on Chicken Stearoyl-CoA desaturase 1
600 gene expression. Biochem. Biophys. Res. Comm. **280**: 25-31.

601

602 Liew, C., Schut, H.A.J., Chin, S.F., Pariza, M.W. and Dashwood R.H. (1995).
603 “Protection of conjugated linoleic acids against 2-amino-3-methylimidazo [4,5-f]
604 quinoline-induced colon carcinogenesis in the f344 rat- a study of inhibitory
605 mechanisms.” Carcinogenesis 16: 3037-3043.

606

607 Martin, G. S., Lunt, D.K., Britian, K.G. and Smith, S.B. 1999. Postnatal
608 development of Stearoyl coenzyme-A desaturase gene expression and adiposity in
609 bovine subcutaneous adipose tissue. J. Anim. Sci. **77**: 630-636.

610

611 Mihara, K. 1990. Structure and regulation of rat liver microsomal Stearoyl Co-A
612 desaturase gene. J. Biochem (Tokyo). **108**: 1022-1029.

613

614 Millar, C.C., Park, Y., Pariza, M.W. and Cook, M.E. 1994. Feeding conjugated
615 linoleic acid to animals partially overcomes catabolic responses to endotoxin
616 injection. Biochem. Biophys.Res. Comm. **198**: 1107-1112.

617

618 Miller, A., Stanton, C. and Devery, R. (2001). “Modulation of arachidonic acid
619 distribution by conjugated linoleic acid isomers and linoleic acid in MCF-7 and
620 SW480 cancer cells.” Lipids 36: 1161-1168.

621

622 Nicolosi, R.J., Courtemanch, K.V., Laitinen, L., Cimeca, J.A. and Huth, P.J. 1993.
623 Effect of feeding diets enriched in conjugated linoleic acid on lipoproteins and aortic
624 atherogenesis in hamsters. Circulation. **88 S 2458**.

625

626 Ntambi, J. M., Buhrow, S.A., Kaestner, K.H., Christy, R.J., Sibley, E., Kelly Jr, T.J.
627 and Lane, M.D. 1988. Differentiation-induced gene expression in 3T3-L1
628 preadipocytes. A second differentially expressed gene encoding Stearoyl-CoA
629 desaturase. *J. Biol. Chem.* **263**: 17291-17300.

630

631 Pariza, M.W., Park, Y. Cook, M., Albright, K. and Liu, W. 1996. Conjugated
632 Linoleic acid (CLA) reduces body fat. *FASEB J.* **10**: 3227.

633

634 Park. Y., Allen, K.G.D. and Shultz, T.D. (2000). "Modulation of MCF-7 breast cancer
635 cell signal transduction by linoleic acid and conjugated linoleic acid in culture."
636 *Anticancer Res.* **20**: 669-676.

637

638 Palombo, J.D., Ganguly, A., Bistran, B.R. and Menard, M.P. (2002). "The
639 antiproliferative effects of biologically active isomers of conjugated linoleic acid on
640 human colorectal and prostatic cancer cells." *Cancer Lett.* **177**: 163-172.

641

642 Peterson, D. G., Kelsey, J.A. and Bauman, D.E. 2002. Analysis of variation in *cis*-9,
643 *trans*-11 conjugated linoleic acid in milk fat of dairy cows. *J. Dairy. Sci.* **85**: 2164-
644 2172.

645

646 Quandt, K., Frech, K., Karas, H., Wingender, E. and Werner, T. (1995). "MatInd and
647 MatInspector-new fast and versatile tools for detection of consensus matches in
648 nucleotide sequence data." *Nucleic Acids Res.* **23**: 4878-4884.

649

650 Shimomura, I., Shimano, H., Korn, B.S., Bashmakov, Y. and Horton, J.D. 1998.
651 Nuclear sterol regulatory element-binding proteins activate genes responsible for the
652 entire program of unsaturated fatty acid biosynthesis in transgenic mouse liver. *J.*
653 *Biol. Chem.* **273**: 35299-35306.

654

655 Solomon, R., Chase, L.E., Ben-Ghedalia, D. and Bauman, D.E. 2000. The effect of
656 non-structural carbohydrate and addition of full fat extruded soybeans on the
657 concentration of conjugated linoleic acid in the milk fat of dairy cows. *J. Dairy Sci.*
658 **83**: 1322-1329.

659

660 Stanton, C., Lawless, F., Kjellmer, G., Harrington, D., Devery, R., Connolly, J.F. and
661 Murphy, J.J. (1997). "Dietary influences on bovine milk cis-9, trans-11 conjugated
662 linoleic acid content." *J. Food. Sci.* **62**: 1083-1086.

663

664 St. John, L.C., Lunt, D.K. and Smith, S.B. 1991. Fatty acid elongation and
665 desaturation enzyme activities of bovine liver and subcutaneous adipose tissue
666 microsomes. *J. Anim. Sci.* **69**: 1064-1073.

667

668 Tabor, D.E., Kim, J.B., Spiegelman, B.M. and Edwards, P.A. 1999. Identification of
669 conserved *cis*-elements and transcription factors required for sterol-regulated
670 transcription of Stearoyl-CoA desaturase 1 and 2. *J. Biol. Chem.* **274**: 20603-20610.

671

672 Visonneau, S., Cesano, A., Tepper, S.A., Scimeca, J.A., Santoli, D. and Kritchevsky,
673 D. (1997). "Conjugated linoleic acid suppresses the growth of human breast
674 adenocarcinoma cells in SCID mice." *Anticancer Res.* **17**: 969-974.

675 Ward, R.J., Travers, M.T., Richards, S., Vernon, R.G., Salter, A.M., Buttery, P.J. and
676 Barber, M.C. 1998. Stearoyl-CoA desaturase mRNA is transcribed from a single
677 gene in the ovine genome. *Biochim. Biophys. Acta.* **1391**: 145-156.
678

679 Waters, K.M., Miller, C.W. and Ntambi, J.M. 1997. Localisation of a
680 polyunsaturated fatty acid response region of Stearoyl-CoA desaturase gene 1.
681 *Biochim. Biophys. Acta.* **1349**:33-42.
682

683 White, S.L., Bertrand, J.A., Wade, M.R., Washburn, S.P., Green Jr, J.T. and Jenkins,
684 T.C. 2001. Comparison of fatty acid content of milk from Jersey and Holstein cows
685 consuming pasture or a total mixed ration. *J. Dairy Sci.* 84:2295-2301.
686

687 Wingender, E., Chen, X., Hehl, R., Karas, H., Liebich, I., Matys, V., Meinhardt, T.,
688 Pruß, M., Reuter, I. and Schacherer, F. (2000). "TRANSFAC: an integrated system
689 for gene expression regulation." *Nucleic Acids Res.* **28**: 316-319.
690

691 Wonsil, B. J., Herbein, J.H. and Watkins, B.A. 1994. Dietary and ruminally derived
692 *trans*-_{18:1} fatty acids alter bovine milk lipids. *J. Nutr.* 124:556-565.
693

694 Xu, J., Nakamura, M.T., Cho, H.P. and Clarke, S.D. 1999. Sterol regulatory element-
695 binding protein-1 expression is suppressed by dietary polyunsaturated fatty acids. A
696 mechanism for the coordinate suppression of lipogenic genes by polyunsaturated fats.
697 *J. Biol. Chem.* **274**:23577-25383.
698

699 Yahagi, N., Shimano, H., Hasty, A.H., Amemiya-Kudo, M., Okazaki, H., Tamura, Y.,
700 Lizuka, Y., Shionoiri, F., Ohashi, K. and Osuga, J. 1999. A crucial role of sterol
701 regulatory element-binding protein-1 in the regulation of lipogenic gene expression by
702 polyunsaturated fatty acids. *J. Biol. Chem.* **274**: 35840-35844.

703

704 Zhang, L., Lan, G.E., Stenn, K. and Prouty, S.M. 2001. Isolation and characterisation
705 of the human Stearoyl-CoA desaturase gene promoter: requirement of a conserved
706 CCAAT *cis*-element. *Biochem. J.* **357**: 183-193.

707

708

709

710

711

712

713

714

715