



Provided by the author(s) and NUI Galway in accordance with publisher policies. Please cite the published version when available.

Title	PPARs and pain
Author(s)	Okine, Bright N.; Gaspar, Jessica C.; Finn, David P.
Publication Date	2018-04-21
Publication Information	Okine, Bright N, Gaspar, Jessica C, & Finn, David P. (2019). PPARs and pain. <i>British Journal of Pharmacology</i> , 176(10), 1421-1442. doi:10.1111/bph.14339
Publisher	Wiley
Link to publisher's version	https://bpspubs.onlinelibrary.wiley.com/doi/10.1111/bph.14339
Item record	http://hdl.handle.net/10379/16256
DOI	http://dx.doi.org/10.1111/bph.14339

Downloaded 2022-05-21T05:51:22Z

Some rights reserved. For more information, please see the item record link above.



REVIEW ARTICLE THEMED ISSUE

PPARs and pain

Correspondence Professor David Finn, Pharmacology and Therapeutics, National University of Ireland Galway, University Road, Galway, Ireland. E-mail: david.finn@nuigalway.ie

Received 13 November 2017; **Revised** 19 February 2018; **Accepted** 26 March 2018

Bright N Okine^{1,2,3,*} , Jessica C Gaspar^{1,2,3,*}  and David P Finn^{1,2,3} 

¹Pharmacology and Therapeutics, National University of Ireland Galway, Galway, Ireland, ²Galway Neuroscience Centre, National University of Ireland Galway, Galway, Ireland, and ³Centre for Pain Research, NCBES, National University of Ireland Galway, Galway, Ireland

*The authors have equal contribution.

Chronic pain is a common cause of disability worldwide and remains a global health and socio-economic challenge. Current analgesics are either ineffective in a significant proportion of patients with chronic pain or associated with significant adverse side effects. The PPARs, a family of nuclear hormone transcription factors, have emerged as important modulators of pain in preclinical studies and therefore a potential therapeutic target for the treatment of pain. Modulation of nociceptive processing by PPARs is likely to involve both transcription-dependent and transcription-independent mechanisms. This review presents a comprehensive overview of preclinical studies investigating the contribution of PPAR signalling to nociceptive processing in animal models of inflammatory and neuropathic pain. We examine current evidence from anatomical, molecular and pharmacological studies demonstrating a role for PPARs in pain control. We also discuss the limited evidence available from relevant clinical studies and identify areas that warrant further research.

Abbreviation

ACC, anterior cingulate cortex; AEA, anandamide; CCI, chronic constriction injury; CFA, complete Freund's adjuvant; DRG, dorsal root ganglion; EMSA, electrophoretic mobility shift assay; FAAH, fatty acid amide hydrolase; iNOS, inducible NOS; OEA, *N*-oleoylethanolamide; PAG, periaqueductal grey; IPAG, lateral PAG; VIPAG, ventrolateral PAG; PEA, *N*-palmitoylethanolamide; PPRE, peroxisome proliferator response element; RXR, retinoid X receptor; RVM, rostroventromedial medulla; SD, Sprague–Dawley; VTA, ventral tegmental area; WKY, Wistar Kyoto 

Peroxisome proliferator-activated receptors

The **PPARs** are ligand-dependent transcription factors that belong to the nuclear hormone superfamily of receptors. Three major isoforms have been identified: **PPAR α** , cloned from mouse liver (Issemann and Green, 1990), **PPAR β/δ** and **PPAR γ** , both cloned from *Xenopus* (Dreyer *et al.*, 1992). These three isoforms share a common structure, typified by the presence of a highly conserved DNA binding domain, with two zinc finger motifs, that recognize the peroxisome proliferator response element (PPRE) in the promoter regions of target genes (Desvergne and Wahli, 1999). They also contain two transcription activation domains; ligand independent AF-1 in the N-terminal domain (Delerive *et al.*, 2002) and the AF-2 in the C-terminal domain, which is ligand-dependent and has a large ligand binding domain. This large ligand binding domain makes it possible for PPARs to interact with a wide array of synthetic and natural lipid ligands.

PPARs exist as heterodimers with the retinoid X receptor (**RXR**), bound to co-repressor proteins in the inactive state. Upon ligand activation, the co-repressors dissociate from the PPAR/RXR complex, allowing for the recruitment of co-activators. The activated PPAR/RXR-co-activator complex subsequently binds to specific DNA sequences or PPRE, resulting in the transcriptional activation of target genes (Green *et al.*, 1992, Tugwood *et al.*, 1992). Genes regulated

by PPARs *via* this PPRE-dependent mechanism are mainly involved in lipid and lipoprotein metabolism (Tugwood *et al.*, 1992). Alternate mechanisms of action that do not involve PPRE binding have been reported, especially for PPAR α , which has known anti-inflammatory effects (Delerive *et al.*, 2001). These latter mechanisms involve inhibition of **NF- κ B** and AP-1 inflammatory signalling and the consequent trans-repression of pro-inflammatory genes such as inducible NOS (**iNOS**), **COX-2** and **TNF- α** (Crisafulli and Cuzzocrea, 2009, Cuzzocrea *et al.*, 2008, Delerive *et al.*, 2000). These anti-inflammatory consequences of PPAR α activation are fundamental to the role of this receptor in modulating both inflammatory and neuropathic pain as discussed later.

PPARs are widely distributed in mammalian tissues, including the peripheral and CNSs (Braissant *et al.*, 1996, Moreno *et al.*, 2004) (Table 1), and are activated by a variety of endogenous compounds derived from saturated and unsaturated fatty acid, of which **N-palmitoylethanolamide** (PEA) (LoVerme *et al.*, 2005) and **N-oleoylethanolamide** (OEA) (Fu *et al.*, 2003) remain the best characterized to date. The ubiquitous distribution of PPARs also reflects their roles in many physiological processes, including an emerging role as key modulators in nociceptive processing. In particular, and as reviewed below, anatomical, molecular and pharmacological studies suggest that the PPAR signalling system may be a viable therapeutic target for the treatment of chronic pain and its comorbidity with stress-related psychiatric disorders.

Table 1

Expression of mRNA or protein for PPAR isoforms within neuroanatomical loci involved in pain

	Neuroanatomical locus	PPAR α	PPAR β/δ	PPAR γ
Peripheral	Dorsal root ganglion	✓ ^{b,c}	?	?
	C fibres	?	?	?
	A δ fibres	?	?	?
	A β fibres	?	?	?
Spinal	Spinal cord	✓ ^{*d}	✓ ^d	✓ ^{a,d,g}
Supraspinal	Frontal cortex	✓ ^d	✓ ^d	✓ ^d
	Pre-frontal cortex (PFC)	✓ ^e	✓ ^e	✓ ^e
	Hippocampus	✓ ^d	✓ ^d	✓ ^d
	Thalamus	✓ ^d	✓ ^d	✓ ^d
	Hypothalamus	X ^d	✓ ^d	✓ ^d
	Basal ganglia	✓ ^{d,e}	✓ ^{d,e}	✓ ^{d,e}
	Amygdala	✓ ^e	✓ ^e	✓ ^e
	PAG	✓ ^f	?	✓ ^f
	Rostroventral medulla (RVM)	?	?	?
	Ventral tegmental area (VTA)	✓ ^e	✓ ^e	✓ ^e

✓, present; X, absent; ?, expression not known.

^aNot expressed in all laminae.

^bD'Agostino *et al.*, 2009.

^cLoVerme *et al.*, 2006.

^dMoreno *et al.*, 2004.

^eWarden *et al.*, 2016.

^fOkine *et al.*, 2017.

^gChuri *et al.*, 2008.

Expression of PPARs in key components of the pain pathway

A role for PPAR signalling in pain processing is suggested by studies demonstrating the presence of the different PPAR isoforms at key peripheral, spinal and supraspinal sites involved in pain processing (Table 1; Figure 1). Within the periphery, PPAR α protein is expressed in the dorsal root ganglion (DRG) (LoVerme *et al.*, 2006), although the relative distribution within DRG sub-nuclei remains unreported. Unlike PPAR α , to our knowledge, the expression of PPAR β or PPAR γ in the DRG

remains unexplored. Despite the paucity of data on the distribution pattern of PPAR α on nociceptive primary afferents (A δ -fibres and C-fibres), the reported analgesic effects of PPAR α agonists, administered locally or peripherally, in animal models of inflammatory and neuropathic pain (see Table 2 for comprehensive list) suggest a modulatory influence on peripheral nociceptive afferents such that PPAR α activation in the DRG results in the suppression or silencing of nociceptive afferent fibre firing. However, the validation of this hypothesis requires further characterization of PPAR α in DRG nuclei using double-labelling immunohistochemistry or *in situ* hybridization techniques.




Figure 1

Anatomical localization of PPAR isoforms in key components of the pain pathway and their role in pain modulation. (1) Okine *et al.* (2014); (2) Okine *et al.* (2017); (3) Okine *et al.* (2016); (4) De Novellis *et al.* (2012); (5) LoVerme *et al.* (2006); (6) Russo *et al.* (2007); (7) Hasegawa-Moriyama *et al.* (2013); (8) Mansouri *et al.* (2017a, b); (9) Churi *et al.* (2008); (10) Griggs *et al.* (2015); (11) Saito *et al.* (2016); (12) Hasegawa-Moriyama *et al.* (2012); (13) Takahashi *et al.* (2011); (14) Sagar *et al.* (2008); (15) D'Agostino *et al.* (2009); (16) Moreno *et al.* (2004); (17) Warden *et al.* (2016); (18) Churi *et al.* (2008); (19) Maeda *et al.* (2008); (20) Chakravarthy *et al.* (2007).

Table 2

Summary of pharmacological studies investigating the role of PPARs in animal models of pain

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Inflammatory	Carrageenan	PEA and GW7647 (PPAR α agonist)	2 μ L per mouse (i.c.v) 0.3 μ L (spinal)	i.c.v and spinal	Swiss mice	Reduced hyperalgesia in mice via inhibition of pro-inflammatory signalling in the carrageenan model of inflammatory pain. The results were mimicked by the PPAR α agonist (GW7647).	D'Agostino <i>et al.</i> (2009)
Inflammatory	Subcutaneous carrageenan	PEA	200, 400 and 800 μ g·mL $^{-1}$	Subcutaneous	Wistar rats	Treatment with PEA reduced allodynia evaluated by Von Frey.	De Filippis <i>et al.</i> (2011)
Inflammatory	Carrageenan	PFOA (PPAR α agonist) and rosiglitazone (PPAR γ agonist)	100 mg·kg $^{-1}$	Systemic	Sprague–Dawley rats	Pretreatment with either drug inhibited carrageenan-induced oedema in a dose-dependent manner and also reduced carrageenan-induced hyperalgesia.	Taylor <i>et al.</i> (2002)
Inflammatory	Carrageenan	Rosiglitazone and 15d-PGJ$_2$ (PPAR γ agonist)	0–5 to 50 mg (rosiglitazone) and 50–200 mg (15d-PGJ $_2$)	i.c.v	Sprague–Dawley rats	i.c.v. administration of the drugs dose-dependently reduced behavioural withdrawal responses to noxious heat. The administration of antagonists (BADCE and GW9662) reversed the anti-hyperalgesic effects.	Morgenweck <i>et al.</i> (2009)
Inflammatory	Carrageenan and collagen-induced arthritis	Adelmidrol (PEA analogue)	10 mg·kg $^{-1}$	Intraperitoneal	Sprague–Dawley rats	The administration of adelmidrol produced a significant inhibition in the development of carrageenan-induced and collagen-induced thermal and mechanical allodynia. This anti-allodynic effect was reversed by GW9662 (PPAR γ antagonist).	Impellizzeri <i>et al.</i> (2016)
Inflammatory	Carrageenan	URB597 (FAAH inhibitor)	25 and 100 μ g	Intraplantar	Sprague–Dawley rats	The administration of URB597 attenuated the hyperalgesia induced by carrageenan. GW6471 (PPAR α antagonist) reversed this effect.	Jhaveri <i>et al.</i> (2008)
Inflammatory	Carrageenan	GW0742 and ATRA (PPAR β/δ agonist)	0.1 mg·kg $^{-1}$ (GW0742) and 5 mg·kg $^{-1}$ (ATRA)	Intraperitoneal	Wistar rats	The administration of both drugs has reduced mechanical and thermal hyperalgesia induced by carrageenan. The co-treatment with a PPAR β/δ antagonist (GSK0660) has blocked the effects of the drugs.	Gill <i>et al.</i> (2013)

continues

Table 2
(Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Inflammatory	Formalin	GW7647 (PPAR α agonist)	0.1–10 μ g/10 μ L	Intraplantar	Swiss mice	GW7647 inhibited phase I and phase II pain behaviour in the formalin-induced inflammatory pain model at the highest dose.	Russo <i>et al.</i> (2007)
Inflammatory	Formalin and carrageenan	PEA	0.01–50 μ g/200 μ L	Intraperitoneal	Swiss mice	Reduction in both early and late phases of formalin-induced nociception by PEA at 5 and 50 mg per paw. PPAR α knockout animals failed to respond to PEA compared to wild-type animals. The injection of carrageenan resulted in a significant reduction of mechanical and thermal threshold values. Both hyperalgesic parameters were strongly reduced by PEA (50 mg).	Sasso <i>et al.</i> (2012)
Inflammatory	Formalin	GW7647 (PPAR α agonist) and GW6471 (PPAR α antagonist)	10 μ g (GW7647); 10 μ g (GW6471)	Intra-mPFC	Sprague–Dawley rats	Intra-mPFC administration of GW6471, but not GW7647, resulted in delayed onset of the early second phase of formalin-evoked nociceptive behaviour. Formalin-evoked nociceptive behaviour was associated with significant reductions in mPFC levels of endogenous PPAR α ligands (PEA and OEA).	Okine <i>et al.</i> (2014)
Inflammatory	Formalin	GW6471 (PPAR α antagonist) and GW9662 (PPAR γ antagonist)	3 nmol–5 μ L (GW6471); 36 nmol–5 μ L (GW9662)	Intra-ACC	Sprague–Dawley rats	Both antagonists significantly reduced formalin-evoked nociceptive behaviour, suggesting facilitatory/permissive roles for these receptors in the ACC in inflammatory pain.	Okine <i>et al.</i> (2017)
Inflammatory	Formalin	GW9662 (PPAR γ antagonist)	14.4 nmol/0.2 μ L	Intra-IPAG	Sprague–Dawley rats and Wistar Kyoto rats	Pharmacological blockade of PPAR γ in the IPAG enhanced formalin-evoked nociceptive behaviour in WKY, but not SD, rats.	Okine <i>et al.</i> (2016)

continues

Table 2
(Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Inflammatory	Formalin and carrageenan	Fenofibrate and pioglitazone	100 or 300 mg·kg ⁻¹ (fenofibrate; acute and chronic – 7 days); 25, 50 or 100 mg·kg ⁻¹ (acute pioglitazone)	<i>Per os</i> (p.o.) – fenofibrate Intraperitoneal – pioglitazone	Swiss mice and Wistar rats	Chronic and acute administration of fenofibrate and acute administration of pioglitazone did not inhibit nociceptive responses of mice in the hot plate or in the first phase of the formalin test. The chronic treatment with fenofibrate and acute administration of pioglitazone (same doses) attenuated the second phase of the formalin-induced nociceptive response. The prolonged treatment with fenofibrate also attenuated the initial phase of the carrageenan-induced nociceptive behaviour in rats.	Oliveira et al. (2007)
Inflammatory	Complete Freund's adjuvant (CFA)-induced inflammation	Rosiglitazone	0.3, 3 or 30 µg	Intraplantar	C57BL/6 mice	Hyperalgesia to mechanical stimuli was dose-dependently attenuated on days 5 and 7 after the procedure in mice that received rosiglitazone, which was reversed to the level of vehicle-injected mice by coadministration of GW9662. In contrast to the effects of rosiglitazone on mechanical stimuli, rosiglitazone had little effect on withdrawal latency to heat stimuli.	Hasegawa-Moriyama et al. (2013)
Inflammatory	Formalin	GW9662 and pioglitazone	2 mg·kg ⁻¹ (GW9662) and 10, 20, 30 or 50 mg·kg ⁻¹ (pioglitazone)	Intraperitoneal	Wistar rats	Pioglitazone at doses 30 and 50 mg·kg ⁻¹ significantly inhibited the flinching behaviour in phase 1 and, at dose 30 mg·kg ⁻¹ , in phase 2. GW9662 had no effect in nociceptive behaviour <i>per se</i> , but it attenuated antinociceptive effects of the combined treatment of simvastatin and pioglitazone.	Mansouri et al. (2017a, b)
Inflammatory	Formalin	GW9662 and pioglitazone	2 mg·kg ⁻¹ (GW9662) and 10, 20, 30 or 50 mg·kg ⁻¹ (pioglitazone)	Intraperitoneal and intraplantar	Wistar rats	Both routes of pioglitazone administration produced antinociception in both phases	Mansouri et al. (2017a, b)

continues

Table 2
(Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Inflammatory	Complete Freund's adjuvant (CFA)-induced	–	50 mg·kg ⁻¹ (pioglitazone)	–	Wistar rats	of formalin-induced pain. Antinociception caused by i.p. and i.pl. pioglitazone was blocked by GW-9662 at doses 2 mg·kg ⁻¹ (i.p.) and 3 µg per paw (i.pl.).	Benani <i>et al.</i> (2004)
Inflammatory	Formalin, carrageenan and writhing test	Oleonic acid (OA)	10, 20 and 40 mg·kg ⁻¹	Oral	Swiss mice	OA treatment inhibits acetic acid-induced abdominal writhing in mice. OA alone did not produce a significant effect on the first phase of the formalin test but reduced the number of paw licks in the second phase of the formalin test.	Vasconcelos <i>et al.</i> (2006)
Inflammatory	Formalin	PEA and GW6471	0.2 and 2 mg·kg ⁻¹ (GW6471 i.p.), 0.2 or 1 g/5 µL per mouse (GW6471 i.t.), 1 µg/20 µL per mouse (GW6471 i.pl.) 1 or 3 mg·kg ⁻¹ (PEA i.p.)	Intraperitoneal (GW6471 was also administered intraplantar and intrathecal)	ICR mice and –/–α7 mice (C57BL/6 background)	GW6471 blocks and PEA potentiates the antinociceptive effects of α7 nAChR full agonist. PEA and GW6471 alone do not affect formalin-evoked nociceptive responses.	Donvito <i>et al.</i> (2017)
Inflammatory	Formalin – Temporomandibular joint (TMJ)	15d-PGJ ₂ and GW9662	0.3, 1 or 3 ng/15 µL per TMJ (GW9662)	Intra-TMJ	Wistar rats	Treatment with 15d-PGJ ₂ attenuated formalin-evoked nociceptive behaviour in the TMJ. This effect was blocked by GW9662 (PPAR _γ antagonist).	Pena-dos-Santos <i>et al.</i> (2009)
Inflammatory	Carrageenan	URBS97, GW6471 and WY14643 (PPAR _α agonist)	25 µg in 50 µL (URBS97) 30 µg in 50 µL (GW6471) 100 µg in 50 µL (WY14643)	Intraplantar	Sprague–Dawley rats	GW6471 completely abolished the inhibitory effects of URBS97 on the carrageenan-evoked expansion of receptive fields (8 g) and WY14643 significantly attenuated carrageenan-evoked expansion of peripheral receptive fields of WDR neurons.	Sagar <i>et al.</i> (2008)
Inflammatory	Carrageenan and formalin –	15d-PGJ ₂ and GW9662	30–300 ng per paw (15d-PGJ ₂)	Intraplantar or intra-TMJ	Wistar rats	15d-PGJ ₂ inhibits the mechanical hypernociception induced carrageenan in the	Napimoga <i>et al.</i> (2007)

continues

Table 2
(Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
	Temporomandibular joint (TMJ)					hind paw and formalin in the TMJ. These effects were blocked by GW9662.	
Inflammatory	Carrageenan	PEA and siRNA for PPAR α	1.0 $\mu\text{g}\cdot\text{h}^{-1}$	i.c.v.	Sprague-Dawley rats	Lower PPAR α expression was observed in the spinal cord of high fat-fed rats. PEA significantly attenuated thermal and mechanical hyperalgesia in HF-fed rats. Intrathecal administration of PPAR α siRNA completely abolished the effects of i.c.v. PEA on pain sensitivity.	Wang et al. (2014)
Inflammatory	Carrageenan (diet-induced obesity)	siRNA for PPAR α	10 μL	Intrathecal	Sprague-Dawley rats	Knockdown of spinal PPAR α eradicated the beneficial effects of ursolic acid on thermal hyperalgesia and paw oedema, and reversed the spinal cord inflammatory response.	Zhang et al. (2016)
Neuropathic and inflammatory	Chronic constriction injury of sciatic nerve (CCI), Freund's adjuvant and carrageenan	PEA and GW7647 (PPAR α agonist)	PEA (50 mg) and GW7647 (50 mg); PEA (20 mg·kg $^{-1}$) s.c.	Intraplantar and subcutaneous	Swiss mice	PEA reduced formalin-induced pain at i.p. doses; GW7647 and PEA reduced hyperalgesic responses in the CCI model of neuropathic pain; acute administration of GW7647 and PEA reduced hyperalgesic responses in the complete Freund's adjuvant and carrageenan models of inflammatory pain.	LoVerme et al. (2006)
Inflammatory and neuropathic	Chronic constriction injury (CCI) and acetic acid-, magnesium sulphate- and kaolin-evoked writhing and formalin	GW9662 and PEA	PEA (20 mg·kg $^{-1}$ per i.p.) GW9662 (2 mg·kg $^{-1}$)	Intraperitoneal	Swiss CD1 mice and PPAR α null type mice	PEA had antihyperalgesic effect on mechanical and thermal stimulus. Single intraperitoneal administration of GW9662 produced a reversion of analgesic effect of both compounds tested (butyrate and FBA).	Russo et al. (2016)
Inflammatory and visceral	Formalin and writhing	OEA WY14643	0.5, 1, 5, 10 and 20 mg·kg $^{-1}$ (OEA)	Intraperitoneal	CD1 mice (wild type and PPAR α null)	Treatment with OEA decreased the writhing response induced	Suardiaz et al. (2007)

continues

Table 2
(Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Neuropathic	Chronic constriction injury of sciatic nerve (CCI)	PEA	5 and 20 mg·kg ⁻¹ (WY14643)	Intraperitoneal	Murine	by acetic acid and formalin-evoked nociceptive behaviour (both phases) in both wild and KO mice. WY14643 did not affect the early phase of the formalin test whereas it slightly decreased the late phase.	Costa <i>et al.</i> (2008)
Neuropathic	Chronic constriction injury of sciatic nerve (CCI)	PEA	10 mg·kg ⁻¹	Intraperitoneal	Murine	Administration of PEA reduced thermal hyperalgesia and mechanical allodynia; these effects were mediated by PPAR _γ .	Costa <i>et al.</i> (2008)
Neuropathic	Chronic constriction injury of sciatic nerve (CCI)	PEA	10 mg·kg ⁻¹	Intraperitoneal for 7 days	C57BL/6j mice	Administration of PEA reduced thermal hyperalgesia and mechanical allodynia.	Bettoni <i>et al.</i> (2013)
Neuropathic	Spared nerve injury	15d-PGJ ₂ and rosiglitazone (PPAR _γ agonists)	25, 50, 100 and 200 μg (15d-PGJ ₂) and 25, 50 and 100 μg (rosiglitazone)	Intrathecal	Sprague-Dawley rats	Treatments with the drugs (dose of 100 μg) decreased mechanical and cold hypersensitivity. Concomitant treatment with PPAR _γ antagonist (BADGE) reversed these effects.	Churi <i>et al.</i> (2008)
Neuropathic	Spared nerve injury	Pioglitazone (PPAR _γ agonist)	1, 3 and 10 mg·kg ⁻¹ ·day ⁻¹ during 7 days (i.p.) 0, 0.3, 3.0, 30.0 mg·kg ⁻¹ daily during 7 days (included in the diet)	Intraperitoneal and oral	Sprague-Dawley RATS	Treatment with pioglitazone had anti-allodynic and anti-hyperalgesic effects, reversed by the PPAR _γ antagonist (GW9662).	Morgenweck <i>et al.</i> (2013)
Neuropathic	Spared nerve injury	Pioglitazone (PPAR _γ agonist)	2 and 10 mg·kg ⁻¹ (i.p.) and 0–300 μg (i.t.)	Intraperitoneal and intrathecal	Sprague-Dawley rats	Treatment rapidly reduced hyperalgesia induced by SNI; administration of GW9662 (PPAR _γ antagonist) reversed the effects.	Griggs <i>et al.</i> (2015)
Neuropathic	Diabetic-induced hyperalgesia	Pioglitazone (PPAR _γ agonist)	30 mg·kg ⁻¹ ·day ⁻¹	Oral (diet)	ZL and ZDF rats	Treatment reduced mechanical and thermal (hot and cold) hyperalgesia in diabetic rats.	Griggs <i>et al.</i> (2016)
Neuropathic	Silk suture thread of the sciatic nerve	Pioglitazone (PPAR _γ agonist)	1–25 mg·kg ⁻¹	Oral (diet)	ICR mice	Treatment attenuated tactile allodynia.	Maeda <i>et al.</i> (2008)
Neuropathic	Partial sciatic nerve ligation	Rosiglitazone (PPAR _γ agonist)	3 and 10 mg·kg ⁻¹	Intraperitoneal and local	C57BL6 mice	Systemic rosiglitazone treatment early in the course of progressive inflammation ameliorated tactile allodynia.	Takahashi <i>et al.</i> (2011)

continues

Table 2
(Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Neuropathic	Diabetic-induced hyperalgesia	PEA	30 mg·kg ⁻¹	Intraperitoneal	Mice	Treatment with PEA relieves mechanical allodynia.	Donvito et al. (2015)
Neuropathic	Paclitaxel-induced allodynia	PEA and GW6471 (PPAR α antagonist)	30 mg·kg ⁻¹ (PEA) and 2 mg·kg ⁻¹ (GW6471)	Intraperitoneal, intraplantar, intrathecal and i.c.v.	ICR mice	PEA treatment had anti-allodynic effects and the treatment with GW6471 (PPAR α antagonist) reversed these effects.	Donvito et al. (2016)
Neuropathic	Chronic constriction injury of sciatic nerve (CCI)	PEA	30 mg·kg ⁻¹ ·day ⁻¹	Subcutaneous (daily)	Wild-type and PPAR α -/- (KO) C57BL6 mice	On day 14, PEA prevented pain threshold alterations in Randall–Selitto and Dynamic Plantar Aesthesiometer tests. In PPAR α null mice PEA treatment failed to induce pain relief.	Di Cesare Mannelli et al. (2013)
Neuropathic	Spared nerve injury	PEA and OEA	10 mg·kg ⁻¹ ·day ⁻¹ during 15 days (OEA and PEA) and 6 nmol per mouse (PEA and OEA)	Intraperitoneal and intra-mPFC	CD-1 mice	Repeated PEA and OEA treatments significantly increased both the thermal and mechanical thresholds in SNI mice. PEA microinjection decreased mechanical threshold with maximum effect at 75 min post-drug. OEA microinjections immediately and transiently reduced mechanical allodynia that lasted up to 30 min post-injection.	Guida et al. (2015)
Neuropathic	Oxaliplatin- induced neuropathy	PEA	30 mg·kg ⁻¹ (acute and chronic administration)	Intraperitoneal	Sprague–Dawley rats	Single administration of PEA was able to reduce oxaliplatin-dependent pain induced by mechanical and thermal stimuli. The repeated treatment with PEA prevented lowering of pain threshold as well as increased pain on suprathreshold stimulation.	Di Cesare Mannelli et al. (2015)
Neuropathic	Spinal cord injury	Pioglitazone	0.5, 1.5 or 3.0 mg·kg ⁻¹ (pioglitazone) and 2 mg·kg ⁻¹ (GW9662)	Intraperitoneal	Sprague–Dawley rats	Pioglitazone treatment significantly increased thermal threshold in spinal cord injured rats compared to the vehicle group. The administration of pioglitazone + GW9662 (PPAR γ antagonist) or GW9662 alone did not result in significant differences to post-SCI surgery rats treated with vehicle.	Park et al. (2007)

continues

Table 2
(Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Neuropathic	Diabetes-induced neuropathic pain	Pioglitazone	10 mg·kg ⁻¹	Oral (chronic administration – 28 days)	Sprague–Dawley rats	The chronic administration of pioglitazone did not attenuate the hyperalgesia induced by the high fat diet/streptozotocin (HFD/STZ) model of diabetes.	Byrne <i>et al.</i> (2015)
Neuropathic	Sciatic nerve ligation (SNL)	Pioglitazone	4.5 and 9.0 mg·mg ⁻¹	Intraperitoneal	Wistar rats	Both doses of pioglitazone attenuated hyperalgesia in the hot plate test and the cold allodynia effect of rats submitted to SNL.	Garg <i>et al.</i> (2017)
Neuropathic	Partial sciatic nerve ligation (PSL)	Pioglitazone	1, 5 or 25 mg·kg ⁻¹	<i>Per os</i> (p.o.) during 5 days	ICR mice	Pioglitazone reduced the tactile allodynia at all doses. However, pioglitazone did not affect nociceptive responses in sham mice.	Iwai <i>et al.</i> (2008)
Neuropathic	Tibial and sural nerve transection (TSNT)	Rosiglitazone	2.5, 5 or 10 mg·kg ⁻¹	<i>Per os</i> (p.o.) daily for 28 days	Wistar rats	Administration of rosiglitazone (at 5 and 10 mg·kg ⁻¹) reduced the mechanical and cold hyperalgesia induced by TSNT without affecting heat hyperalgesia.	Jain <i>et al.</i> (2009)
Neuropathic	Lumbar 5 spinal nerve transection	Pioglitazone	2.5, 5 or 10 mg·kg ⁻¹	<i>Per os</i> (p.o.) daily for 14 days	Sprague–Dawley rats	Pioglitazone (5 and 10 mg·kg ⁻¹) attenuated mechanical hyperalgesia produced by lumbar 5 spinal nerve transection	Jia <i>et al.</i> (2010)
Neuropathic	Trigeminal inflammatory compression (TIC)	Pioglitazone, GW0742 (PPAR β/δ agonist), bezafibrate, fenofibrate, GW9662	100, 300 or 600 mg·kg ⁻¹ (pioglitazone); 1 or 6 mg·kg ⁻¹ (GW0742); 100 mg·kg ⁻¹ (bezafibrate); 200 mg·kg ⁻¹ (fenofibrate); 30 mg·kg ⁻¹ (GW9662)	Oral – pioglitazone 600 mg·kg ⁻¹ and bezafibrate 100 mg·kg ⁻¹ Intraperitoneal – all the others	C57BL/6 mice	Systemic administration of pioglitazone attenuates whisker pad mechanical allodynia at doses of 300 and 600 mg·kg ⁻¹ . Administration of GW9662 prior to pioglitazone (300 mg·kg ⁻¹) blocked the analgesic effect of pioglitazone. GW0742 (6 mg·kg ⁻¹) partially attenuated mechanical allodynia in mice with TIC injury compared to vehicle treated mice.	Lyons <i>et al.</i> (2017, 2018)
Neuropathic	Chronic constriction injury (CCI)	Pioglitazone	20 mg·kg ⁻¹ ·day ⁻¹	Oral – daily for 14 days	Sprague–Dawley rats	Pioglitazone attenuated the CCI-induced mechanical and thermal hyperalgesia.	Murad and Ayuob (2015)

continues

Table 2
 (Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Neuropathic	Spinal nerve ligation (SNL)	Pioglitazone	5, 10 or 20 mg·kg ⁻¹	Intraperitoneal – daily for 28 days	Sprague–Dawley rats	Higher doses of pioglitazone attenuated the SNL-induced mechanical allodynia, cold allodynia, mechanical hyperalgesia and thermal hyperalgesia.	Pottabathini et al. (2016)
Neuropathic	Oxaliplatin-induced neuropathic pain	15 Thiazolidinones (TZDs) and GW9662	40 mg·kg ⁻¹ (TZDs) 4 mg·kg ⁻¹ (GW9662)	Intraperitoneal	C57BL/6 mice	Except for compound 14, all TZDs showed antinociceptive properties; these TZDs attenuated oxaliplatin-induced mechanical hyperalgesia. This effect was prevented by GW9662 (PPAR γ antagonist).	Moreira et al. (2017)
Neuropathic	Sciatic nerve ligation	GW6471	5, 10 or 20 mg·kg ⁻¹	Intraperitoneal	C57BL/6 wild-type and PPAR α null lineages	Higher sensitivity to thermal and mechanical non-noxious and noxious stimuli, and cold and mechanical allodynia and heat hyperalgesia was observed in mice lacking PPAR α . Writting after acetic acid were also enhanced in mutant mice. The blockade of PPAR α did not alter nociceptive behaviour.	Ruiz-Medina et al. (2012)
Neuropathic	Oxaliplatin-induced neuropathic pain	Rosiglitazone	3 and 10 mg·kg ⁻¹	Per os (p.o.) daily for 20 days	Sprague–Dawley rats	Rosiglitazone attenuated hyperalgesia and allodynia resulted by oxaliplatin neuropathy.	Zanardelli et al. (2014)
Neuropathic	Trigeminal nerve injury (TNI)	Pioglitazone	100 mg·kg ⁻¹	Intraperitoneally – daily for 7 days	C57BL/6 mice	DCS (NMDA agonist) and pioglitazone combination n attenuated orofacial neuropathic pain and anxiety related behaviours. The treatment with pioglitazone alone did not alter nociceptive behaviour.	Lyons et al. (2017, 2018)
Neuropathic	Peripheral nerve injury – L5 Spinal nerve transection	Pioglitazone and GW9662	10 mg·kg ⁻¹ (pioglitazone) 2 mg·kg ⁻¹ (GW9662)	Intraperitoneal	Sprague–Dawley rats	Pioglitazone improved the mechanical hyperalgesia in operated rats. This effect was reversed by GW9662.	Jia et al. (2013)
Neuropathic	Chronic constriction injury (CCI)	Pioglitazone and GW9662	10 mg·kg ⁻¹ (pioglitazone) 2 mg·kg ⁻¹ (GW9662)	Intraperitoneal	Sprague–Dawley rats	Treatment with pioglitazone attenuated CCI-induced hyperalgesia. This effect was reversed by GW9662 administration.	Jia et al. (2016)

continues

Table 2
(Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Neuropathic and Visceral	Acetic acid induced visceral pain, sciatic nerve injury (SNI)	MK886 (PPAR α antagonist)	2 mg·kg ⁻¹	Intraperitoneal	ICR mice, Kunming mice and C57BL/6 mice and PPAR- α knockout mice (-/-)	F96 (selective NAAA inhibitor) had an overall antinociceptive effect in the different models and tests carried out in the study. This effect was widely blocked by PPAR α antagonist MK886 and by genetic disruption of PPAR- α .	Yang <i>et al.</i> (2015)
Visceral	Diarrhoea-predominant irritable bowel syndrome (D-IBS)	Pioglitazone and GW9662	2 mg·kg ⁻¹ (pioglitazone); 3 mg·kg ⁻¹ (GW9662)	Intraperitoneal on days 7, 9 and 11	Wistar rats	Pioglitazone reduced visceral hypersensitivity and defecation frequency and increased nociceptive thresholds.	Paragomi <i>et al.</i> (2014)
Other	-	PEA	3 and 6 nmol	Intra-vIPAG	Wistar rats	Reduced thermo-nociceptive threshold, as well as on/off cell activity in the rostro-ventromedial medulla (RVM).	De Novellis <i>et al.</i> (2012)
Other	-	Pioglitazone and GW9662 (PPAR γ antagonist)	10 or 30 mg·kg ⁻¹ (pioglitazone) 2.5 or 5 mg·kg ⁻¹ (GW9662)	Oral (via gavage) – Daily (concomitant with morphine administration or for the prior 2 days before testing)	C57 mice	Treatment with pioglitazone attenuated the development of morphine tolerance. Pioglitazone administration in mice that were not chronically treated with morphine does not have an effect in nociception. Pretreatment with GW9662 reversed the effects of pioglitazone in morphine-treated rats. GW9662 alone does not have an effect in nociceptive responses. The development of tolerance for morphine is more pronounced in PPAR γ knockout mice.	De Guglielmo <i>et al.</i> (2014)
Other	-	Pioglitazone and GW9662	20 or 40 mg·kg ⁻¹ (pioglitazone) 2 mg·kg ⁻¹ (GW9662)	Per os (p.o.) (pioglitazone) – daily for 17 days concomitant with morphine treatment Subcutaneous (GW9662)	Wistar rats	Treatment with pioglitazone attenuated the development of morphine tolerance. GW9662 administration 30 min before pioglitazone antagonized the mentioned pioglitazone-induced effects.	Chavimi <i>et al.</i> (2015)

continues

Table 2
 (Continued)

Type of pain	Model	Drug	Dose	Route of administration	Animal	Outcome	Reference
Other	-	Pioglitazone and GW9662	5, 10, 20 or 40 mg·kg ⁻¹ (pioglitazone) 2 mg·kg ⁻¹ (GW9662)	Per os (p.o.) (pioglitazone) – daily for 17 days concomitant with morphine treatment Subcutaneous (GW9662) – daily before pioglitazone administration	Wistar Rats	The highest dose of pioglitazone <i>per se</i> did not alter the pain threshold in tail-flick test. Treatment with pioglitazone attenuated the development of morphine tolerance and GW-9662 administration 30 min before pioglitazone antagonized the pioglitazone-induced effects.	Chavimi <i>et al.</i> (2014)
Other	Incisional pain	Rosiglitazone	25 µg	Intraplantar (in loci)	BKS.Cg – +Leprdb/ +Leprdb/Jcl Mice	Rosiglitazone alleviates mechanical hyperalgesia resulted by the incision.	Saito <i>et al.</i> (2016)
Other	Incisional pain	Rosiglitazone	0.5 mg·mL ⁻¹	Local (intraplantar)	C57BL/6 mice	Local administration of rosiglitazone immediately after the procedure ameliorates thermal and mechanical hyperalgesia.	Hasegawa-Moriyama <i>et al.</i> (2012)

The Table was constructed based on searches within Pubmed central (<https://www.ncbi.nlm.nih.gov/pmc/>) and Google scholar (<https://scholar.google.co.uk/>) using combinations of one or more of the following search terms: PPARs, PPAR α , PPAR β , PPAR γ , PEA, OEA, synthetic PPAR ligands, pharmacological effects, CNS, brain, spinal cord, dorsal root ganglion, animal models and pain.

PPAR α protein expression in the spinal cord has also been demonstrated in earlier studies (Benani *et al.*, 2004, Okine *et al.*, 2015). The functional relevance of PPAR α signalling in the spinal cord to nociceptive processing is demonstrated at least in part by the reported increases in PPAR α activation or protein expression in animal models of chronic inflammatory and neuropathic pain states. For example, using electrophoretic mobility shift assays (EMSA), Benani and colleagues were able to demonstrate a rapid increase in activation of the PPAR α protein isoform in the rat spinal cord after injection of complete Freund's adjuvant into the hind paw (Benani *et al.*, 2004). Moreover, increased PPAR α protein expression in the ipsilateral spinal cord was observed in the rat spinal nerve ligation model of neuropathic pain (Okine *et al.*, 2015). Furthermore, down-regulation of PPAR α protein expression in the spinal cord contributed to augmented peripheral inflammation and inflammatory hyperalgesia in diet-induced obese rats (Wang *et al.*, 2014). While the pathophysiological relevance of PPAR α activation or changes in PPAR α protein expression in the spinal cord during hyperalgesia requires further investigation, these findings provide evidence for PPAR α , as a potentially important player in spinal pain processing. In addition to PPAR α , PPAR γ is also expressed in the spinal cord (Table 1). Increased PPAR γ protein expression in the spinal trigeminal caudalis 3 weeks after trigeminal nerve injury in mice has been reported to play a significant role in trigeminal nociceptive transmission, as demonstrated by the attenuation of whisker pad mechanical allodynia (Lyons *et al.*, 2017), and identifies PPAR γ as a potential therapeutic target for orofacial neuropathic pain.

In comparison, there is a paucity of data on the expression or activation of PPAR β/δ in the spinal cord in inflammatory or neuropathic pain states. In this regard, further characterization of PPAR β/δ is essential to establish whether changes in expression or endogenous activation of the receptor occur in pain states, and the extent to which they contribute to spinal pain processing.

As shown in Table 1, all three PPAR isoforms are widely expressed supraspinally, in key brain regions involved in pain processing. The expression of PPARs at key relay sites such as the thalamus and the midbrain periaqueductal grey (PAG) may reflect a role for PPAR signalling in modulating the activity of ascending and descending pain pathways. Furthermore, the presence of PPARs within cortical regions and the amygdala suggests potential involvement of PPAR signalling in modulating cognitive or affective components of pain. While there is currently no direct evidence in support of this hypothesis, this view is however consistent with the role of both the cortex and amygdala as key brain regions involved in the modulation of the cognitive-affective components of pain (see Fuchs *et al.*, 2014; Neugebauer, 2015).

Evidence from pharmacological or genetic manipulation studies for a role of PPARs in pain

Pharmacological or genetic manipulation of PPAR α and PPAR γ using selective agonists, antagonists or gene knockout

approaches specifically targeting these receptors within the pain pathway has been shown to alter nociceptive processing, demonstrated by changes in neuronal activity or behavioural responses in animal models of inflammatory and neuropathic pain (Figure 1). Both PPAR α and PPAR γ regulate the release of pro-inflammatory mediators associated with tissue or nerve injury through the inhibition of pro-inflammatory signalling pathways such as NF- κ B activation (Cuzzocrea *et al.*, 2008, Delerive *et al.*, 2000) and suppression of downstream pro-inflammatory molecules including COX-2 and iNOS (D'Agostino *et al.*, 2009), two key players in the development of chronic pain states. The majority of pharmacological studies to date demonstrate antinociceptive effects of both endogenous and synthetic agonists of PPAR α and PPAR γ in animal models of inflammatory and neuropathic pain (for a comprehensive summary of the published studies to date see, Table 2). It is pertinent to note that a significant proportion of preclinical studies investigating the role of endogenous PPAR ligands in nociceptive processing in the periphery and the CNS have mainly focused on the effects of PEA, with relatively little attention given to the role of OEA. One possible reason that may account for this apparent bias is the reported activation of **TRPV1** pronociceptive non-selective cation channels (the vanilloid receptors), by OEA (Ahern, 2003). Thus, it is possible that any PPAR-mediated analgesic effects of OEA are likely to be nullified by its TRPV1-mediated pronociceptive effects, as demonstrated in an animal model of neuropathic pain (Guida *et al.*, 2015).

The pharmacological effects of PEA involve both transcription-dependent and transcription-independent or non-genomic mechanisms. While the former account primarily for the anti-inflammatory effects associated with PPAR activation, the non-genomic mechanisms are thought to underlie the rapid antinociceptive effects of not only PEA but also other synthetic PPAR agonists in animal models of inflammatory and neuropathic pain (Churi *et al.*, 2008, LoVerme *et al.*, 2006; Donvito *et al.*, 2015). It is, however, important to note that the non-genomic mechanisms mediating the effects of PEA are not independent of PPAR expression or activation. Indeed, evidence from studies with PPAR knockout mice suggests that the modulation of medium and large **Ca²⁺ channels** associated with the rapid antinociceptive effects of PEA and other synthetic PPAR α agonists on inflammatory pain behaviour in mice are contingent upon PPAR α receptor expression in the DRG (LoVerme *et al.*, 2006). Given that these rapid antinociceptive effects are incompatible with the duration of longer term transcription-dependent mechanisms, the modulation of Ca²⁺ channels in this instance may be a by-product of protein-protein interactions induced by changes in PPAR protein conformation following the binding of agonist to the receptor. Further studies aimed at identifying the precise mechanisms involved could potentially lead to the identification of novel therapeutic targets for the treatment of pain.

The non-genomic effects of PEA may also involve the indirect activation of other receptors, such as cannabinoid receptors, mediated by anandamide (**AEA**) through the so-called 'entourage hypothesis' (LoVerme *et al.*, 2005). In this context, competition by PEA for fatty acid amide hydrolase (**FAAH**)-mediated hydrolysis, is thought to provide a

'sparing effect' on AEA hydrolysis by FAAH, resulting in enhanced signalling at endocannabinoid targets, in particular **CB₁** or **CB₂** receptors, to produce analgesia (Tuo *et al.*, 2017). In keeping with this, we have recently demonstrated a role for CB₁ receptors in the antinociceptive effects of PEA injected directly into the anterior cingulate cortex (ACC) in the rat formalin test (Okine *et al.*, 2016). Moreover, given the preferential binding of AEA over PEA to PPAR γ (Bouaboula *et al.*, 2005), it is possible that entourage-mediated signalling involving AEA is likely to underpin the PPAR γ -mediated antinociceptive effects of PEA (Costa *et al.*, 2008). Indeed, AEA binds to and activates PPAR α in addition to PPAR γ (O'Sullivan and Kendall, 2010). Here, it is important to note that these findings have implications not only for the interpretation of data demonstrating the analgesic effects of endogenous PPAR ligands such as PEA and OEA or inhibitors of FAAH which enhance levels of endogenous PPAR agonists but also the design of such experiments given the possible involvement of other receptor systems other than PPARs. In this regard, the inclusion of appropriate robust control experiments involving PPAR antagonists or gene knockout should always be considered when examining a role for PPARs as mediators of the pharmacological effects of endogenous PPAR ligands.

The analgesic effects of PPAR agonists may also be mediated *via* modulation of cellular organelles. For example, a combination drug therapy of the synthetic PPAR γ agonist **pioglitazone** with **D-cycloserine** attenuates chronic orofacial neuropathic pain and associated anxiety by improving mitochondrial function following trigeminal nerve injury (Lyons *et al.*, 2018). With the molecular mechanisms continuing to be elucidated, these findings widen the scope and increase the appeal of PPAR agonists as therapeutic agents for treating pain. Furthermore, given the involvement of both genomic and non-genomic mechanisms in mediating the effects of PPAR agonists, further studies aimed at determining which mechanisms are predominant in different types of pain will be important for the optimization of the analgesic effects of PPAR agonists.

It is however important to note that while the weight of evidence is in favour of antinociceptive effects of PPAR α or PPAR γ activation at multiple sites within the pain pathway, recent findings from our group also reveal a pain permissive or facilitatory role for PPAR signalling in discrete brain regions such as the ACC (Okine *et al.*, 2017, Okine *et al.*, 2014). Intra-ACC injection of **GW6471** (selective PPAR α antagonist) or **GW9662** (selective PPAR γ antagonist) significantly suppressed the onset of formalin-evoked nociceptive behaviour in rats (Okine *et al.*, 2017). Such permissive or facilitatory roles of endogenous PPAR activation within the ACC may allow the animal to perceive pain and take the necessary actions to escape from, or avoid injury.

The specific role of PPAR β/δ activation in pain processing remains largely unknown, despite molecular evidence demonstrating the presence of the receptor at key sites within the pain pathway such as the spinal cord, thalamus and PAG (Table 1; Figure 1). However, in a previous study, administration of **GW0742**, a selective PPAR β/δ receptor agonist (0.1 mg·kg⁻¹ per i.p. for 4 days) significantly decreased mechanical and thermal hyperalgesia in adult male Wistar rats, induced by carrageenan injection into the hind paw,

compared with vehicle-treated controls. These effects were reversed in the presence of the selective PPAR β/δ antagonist **GSK0660** (0.3 mg·kg⁻¹ per i.p. for 4 days) (Gill *et al.*, 2013). These findings demonstrate the potential of PPAR β/δ agonists as therapeutic agents for the treatment pain. Further preclinical studies are however needed to understand fully the extent to which PPAR β/δ -mediated signalling modulates nociceptive transmission within the CNS.

Evidence from clinical trials

Over the last couple of decades, the analgesic effects of PEA, an endogenous PPAR agonist, or its derivatives, have been demonstrated in several clinical trials in different pain conditions. In a recent comprehensive review of 21 clinical trials, Gabriellsson *et al.*, reported that oral or sublingual treatment with PEA or micronized PEA (PEA- μ m; reduced crystal particles of PEA that enhance the dissolution and reduce the absorption variability) reduced pain intensity in patients with neuropathic and inflammatory joint pain phenotypes (Gabriellsson *et al.*, 2016). Significantly, these treatments were not associated with any marked side effects. Similar reports of the analgesic effects of PEA in clinical trials have been discussed in another comprehensive review by Hesselink and Hekker (2012). These studies report that administration of PEA (doses ranging from 300 to 600 mg·day⁻¹; mostly orally administered as tablets) is effective against a range of pain conditions including neuropathic pain, low back pain and postoperative pain.

In contrast, Andresen and colleagues report that a 12-week treatment with PEA- μ m did not alleviate pain in patients with spinal cord injury-induced neuropathic pain, compared to placebo-treated patients (Andresen *et al.*, 2016). The authors however point out that the limited knowledge on PEA- μ m pharmacokinetics, including information on diffusion to the CSF, make it difficult to draw more specific conclusions. It is also possible that the heterogeneity in the population of spinal cord injury pain phenotypes could have affected the outcome of this study. These clinical effects of PEA however, while suggestive of a role for PPAR signalling, do not necessarily rule out the involvement of other receptor systems, given the multiple signalling pathways mediating the pharmacological effects of PEA as demonstrated in preclinical studies and discussed elsewhere in this review. In this regard, the use of synthetic PPAR agonists in clinical trials may be more beneficial and informative. In keeping with this line of argument, a more defined role for PPAR signalling in modulating human pain conditions was demonstrated in a study by Smith and colleagues, who reported a reduction in occurrence of myalgia, a musculoskeletal pain disorder, in men receiving **clofibrate**, an approved PPAR α agonist used clinically for the treatment of dyslipidemia (Smith *et al.*, 1970). However, subsequent attempts at replicating these early promising results using other fibrates to alleviate muscular pain have not been successful (Biga *et al.*, 2005). Nevertheless, these drugs were found to be effective in attenuating pain associated with rheumatoid arthritis and osteoarthritic pain (van Eekeren *et al.*, 2013). These findings indicate that synthetic PPAR α agonists can have analgesic effects in specific types of pain.

Synthetic agonists of PPAR γ are currently used clinically as insulin sensitizers in the treatment of non-insulin dependent (Type 2) diabetes. However, despite preclinical evidence demonstrating their analgesic effects in a variety of animal models of inflammatory and neuropathic pain (see Table 2), to our knowledge, there are currently no published clinical studies investigating their effects on pain in human subjects or patients. Similarly, there is a paucity of clinical studies investigating the effects of synthetic PPAR β agonists on pain.

A potential role for PPAR signalling in interactions between pain and negative affect

The close relationship between stress (and stress-related disorders such as anxiety and depression) and chronic pain is now widely recognized (Jennings *et al.*, 2014, Olango and Finn, 2014). Although the role of PPAR signalling in the modulation of stress-pain interactions remains largely unexplored, the abundant expression of PPARs in key brain regions involved in the modulation of the cognitive and affective components of pain such as the amygdala and PAG (Table 1 and Figure 1), and the availability of endogenous PPAR ligands in these brain structures, supports a potential role for the PPAR signalling system in stress-pain interactions and as a potential therapeutic target for the treatment of comorbid chronic pain and affective disorders. This view is also consistent with our recent demonstrations of enhanced second phase formalin-evoked nociceptive behaviour following selective blockade of PPAR γ in the lateral PAG in Wistar Kyoto (WKY) but not Sprague-Dawley (SD) rats (Okine *et al.*, 2017). The WKY rat strain is stress-hyperresponsive and exhibits a hyperalgesic phenotype to nociceptive stimuli compared with SD rats and is considered a suitable genetic model for studying stress-pain interactions (Burke *et al.*, 2010, Rea *et al.*, 2014, O'Mahony *et al.*, 2013). While the specific contribution of PPAR γ signalling to the stress hyperresponsive

phenotype of WKY rats is currently not clear, the differential effects of pharmacological modulation of PPAR γ in the lateral PAG on formalin-evoked nociceptive behaviour in SD and WKY rats suggests an important role for this receptor in a genetic background that is prone to stress and hypersensitivity to nociceptive stimuli. These findings also suggest that PPAR γ -mediated signalling in the lateral PAG may represent a potential therapeutic target for future development of effective therapies for treating comorbid chronic pain and stress-related disorders such as anxiety and depression. The therapeutic potential of PPAR γ for treatment of pain and mood disorder comorbidity is also supported by evidence that pioglitazone attenuates chronic constriction injury (CCI)-induced depression-related behaviour in the forced swim test in rats (Garg *et al.*, 2017)), reduces anxiety-like behaviour in a mouse model of chronic orofacial neuropathic pain (Lyons *et al.*, 2018) and augments both the antidepressant and the antinociceptive effects of **fluoxetine** in the rat CCI model of neuropathic pain (Murad and Ayuob, 2015). Additional studies on the therapeutic potential of PPAR agonists (including those for PPAR α and PPAR β/δ) for treatment of the affective/emotional component of chronic pain are warranted.

Future perspectives

Synthetic PPAR α and PPAR γ agonists such as clofibrate and **rosiglitazone**, respectively, are currently available clinically for the treatment of medical conditions such as dyslipidemia and non-insulin dependent (Type 2) diabetes, the latter of which is often associated with altered sensory processing in the extremities. The current clinical use of these drugs provides an opportunity for clinical trials to explore and evaluate their potential as novel analgesics for the treatment of intractable chronic pain conditions such as diabetic neuropathy.

It is however important to note that the development of analgesics targeting the PPAR signalling system must also take into account their potential adverse side effects

Table 3

Summary of adverse effects associated with PPAR activation

Cardiovascular effects	• Increased risk of myocardial infarction ^{a,b,c}
Tumorigenic effects	• Increased risk of bladder cancer in patients on long-term therapy with synthetic PPAR agonists ^{c,d}
Others peripheral effects	• Haemangioma and renal pelvic tumours ^d
	• PPAR agonist induced peripheral oedema formation <i>via</i> fluid retention ^{e,f}
	• Increased liver toxicity ^g
	• Inhibition of gastrointestinal motility ^h
	• Inhibition of human aortic smooth-muscle contraction associated with PPAR α activation ⁱ

^a(Nissen and Wolski, 2007).

^b(Loke *et al.*, 2011).

^c(Friedland *et al.*, 2012).

^d(Aoki, 2007).

^e(Sunder Mudaliar *et al.*, 2003).

^f(Beltowski *et al.*, 2013).

^g(Floyd *et al.*, 2009).

^h(Capasso *et al.*, 2001).

ⁱ(Staels *et al.*, 1998).

(Table 3), given the important physiological roles of the PPAR signalling system. For example, the inhibitory effects of PEA on gastrointestinal motility (Capasso *et al.*, 2001) or on contraction of human aortic smooth muscle, associated with PPAR α activation (Staels *et al.*, 1998,) may limit the therapeutic efficacy of PPAR-targeting analgesics. Nonetheless, evidence from clinical studies using PEA so far suggests that these side effects are unlikely to be of major concern over a short duration of administration of PPAR agonist and that the therapeutic benefits may outweigh the side effects. Moreover, the demonstration of cardioprotective and anti-tumour effects (Wright *et al.*, 2014) of some synthetic PPAR agonists, but not others, suggests that these reported adverse side effects may be agonist-dependent rather than PPAR target-dependent. Further studies are however required to determine the long-term effects of persistent pharmacological modulation of PPAR signalling in chronic pain states.

The role of PPAR signalling in the development or modulation of other chronic pain conditions, such as osteoarthritis, cancer pain and migraine requires further study, as does the interaction of PPAR signalling with other well-characterized endogenous pain control systems and currently prescribed analgesics. On this latter point, the PPAR γ agonist pioglitazone has been shown to attenuate tolerance to **morphine** in a rat model of inflammatory pain (Ghavimi *et al.*, 2015, Ghavimi *et al.*, 2014) and in the mouse tail immersion test (de Guglielmo *et al.*, 2014). Similar potential synergistic antinociceptive interactions with the cannabinoid (Russo *et al.*, 2007) and TRPV1 (Ambrosino *et al.*, 2014) signalling systems have been reported. In respect of the latter study, the evidence suggests that the potential antinociceptive effects of this synergistic interaction are likely to be facilitated by PPAR α -dependent activation of TRPV1 channels and subsequent desensitization of the receptor. Elsewhere, a synergistic antinociceptive interaction between PEA and the opiate drug, **tramadol**, has been demonstrated in the mouse formalin test (Déciga-Campos *et al.*, 2015). The antinociceptive mechanisms of the PEA and tramadol combination involved the opioid receptor, TRPV1 and PPAR α . Significantly, the sedative effects of the combination of PEA and tramadol were minimal compared with those observed with individual treatments. Collectively, these findings make a compelling case for an increased understanding of PPAR signalling and its crosstalk with other analgesic targets. Such knowledge could lead to the development of novel PPAR signalling based-analgesic strategies.

Nomenclature of targets and ligands

Key protein targets and ligands in this article are hyperlinked to corresponding entries in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS Guide to PHARMACOLOGY (Harding *et al.*, 2018), and are permanently archived in the Concise Guide to PHARMACOLOGY 2017/18 (Alexander *et al.*, 2017a,b,c).

Acknowledgements

This work was funded by grants from Science Foundation Ireland (10/IN.1/B2976), the Irish Research Council, CNPq -

Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil (207530/2014-9) and the National University of Ireland Galway.

Conflict of interest

The authors declare no conflicts of interest.

References

- Ahern GP (2003). Activation of TRPV1 by the satiety factor oleoylethanolamide. *J Biol Chem* 278: 30429–30434.
- Alexander SPH, Kelly E, Marrion NV, Peters JA, Faccenda E, Harding SD *et al.* (2017a). The Concise Guide to PHARMACOLOGY 2017/18: Nuclear hormone receptors. *Br J Pharmacol* 174: S208–S224.
- Alexander SPH, Kelly E, Marrion NV, Peters JA, Faccenda E, Harding SD *et al.* (2017b). The Concise Guide to PHARMACOLOGY 2017/18: Voltage-gated ion channels. *Br J Pharmacol* 174: S160–S194.
- Alexander SPH, Christopoulos A, Davenport AP, Kelly E, Marrion NV, Peters JA *et al.* (2017c). The Concise Guide to PHARMACOLOGY 2017/18: G protein-coupled receptors. *Br J Pharmacol* 174: S17–S129.
- Ambrosino P, Soldovieri MV, De Maria M, Russo C, Tagliatela M (2014). Functional and biochemical interaction between PPAR α receptors and TRPV1 channels: potential role in PPAR α agonists-mediated analgesia. *Pharmacol Res* 87: 113–122.
- Andresen SR, Bing J, Hansen RM, Biering-Sorensen F, Johannesen IL, Hagen EM *et al.* (2016). Ultramicrosized palmitoylethanolamide in spinal cord injury neuropathic pain: a randomized, double-blind, placebo-controlled trial. *Pain* 157: 2097–2103.
- Aoki T (2007). Current status of carcinogenicity assessment of peroxisome proliferator-activated receptor agonists by the US FDA and a mode-of-action approach to the carcinogenic potential. *J Toxicol Pathol* 20: 197–202.
- Beltowski J, Rachańczyk J, Włodarczyk M (2013). Thiazolidinedione-induced fluid retention: recent insights into the molecular mechanisms. *PPAR Res* 2013: 628628.
- Benani A, Heurtaux T, Netter P, Minn A (2004). Activation of peroxisome proliferator-activated receptor alpha in rat spinal cord after peripheral noxious stimulation. *Neurosci Lett* 369: 59–63.
- Bettoni I, Comelli F, Colombo A, Bonfanti P, Costa B (2013). Non-neuronal cell modulation relieves neuropathic pain: efficacy of the endogenous lipid palmitoylethanolamide. *CNS Neurol Disord Drug Targets* 12: 34–44.
- Biga J, Vigreux P, Lapeyre-Mestre M, Montastruc JL (2005). Muscular pain and hypolipidaemic drugs: a cross-sectional study in cardiological outpatient practice. *Therapie* 60: 295–297.
- Bouaboula M, Hilairat S, Marchand J, Fajas L, Fur GL, Casellas P (2005). Anandamide induced PPAR γ transcriptional activation and 3T3-L1 preadipocyte differentiation. *Eur J Pharmacol* 517: 174–181.
- Braissant O, Fougelle F, Scotto C, Dauca M, Wahli W (1996). Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR-alpha, -beta, and -gamma in the adult rat. *Endocrinology* 137: 354–366.
- Burke NN, Hayes E, Calpin P, Kerr DM, Moriarty O, Finn DP *et al.* (2010). Enhanced nociceptive responding in two rat models of

- depression is associated with alterations in monoamine levels in discrete brain regions. *Neuroscience* 171: 1300–1313.
- Byrne FM, Cheetham S, Vickers S, Chapman V (2015). Characterisation of pain responses in the high fat diet/streptozotocin model of diabetes and the analgesic effects of antidiabetic treatments. *J Diabetes Res* 2015: 752481.
- Capasso R, Izzo AA, Fezza F, Pinto A, Capasso F, Mascolo N *et al.* (2001). Inhibitory effect of palmitoylethanolamide on gastrointestinal motility in mice. *Br J Pharmacol* 134: 945–950.
- Chakravarthy MV, Zhu Y, López M, Yin L, Wozniak DF, Coleman T *et al.* (2007). Brain fatty acid synthase activates PPAR α to maintain energy homeostasis. *J Clin Invest* 117: 2539–2552.
- Churi SB, Abdel-Aleem OS, Tumber KK, Scuderi-Porter H, Taylor BK (2008). Intrathecal rosiglitazone acts at peroxisome proliferator-activated receptor-gamma to rapidly inhibit neuropathic pain in rats. *J Pain* 9: 639–649.
- Costa B, Comelli F, Bettoni I, Colleoni M, Giagnoni G (2008). The endogenous fatty acid amide, palmitoylethanolamide, has anti-allodynic and anti-hyperalgesic effects in a murine model of neuropathic pain: involvement of CB(1), TRPV1 and PPAR γ receptors and neurotrophic factors. *Pain* 139: 541–550.
- Crisafulli C, Cuzzocrea S (2009). The role of endogenous and exogenous ligands for the peroxisome proliferator-activated receptor alpha (PPAR-alpha) in the regulation of inflammation in macrophages. *Shock* 32: 62–73.
- Cuzzocrea S, Bruscoli S, Mazon E, Crisafulli C, Donato V, Di Paola R *et al.* (2008). Peroxisome proliferator-activated receptor- α contributes to the anti-inflammatory activity of glucocorticoids. *Mol Pharmacol* 73: 323–337.
- D'Agostino G, La Rana G, Russo R, Sasso O, Iacono A, Esposito E *et al.* (2009). Central administration of palmitoylethanolamide reduces hyperalgesia in mice via inhibition of NF- κ B nuclear signalling in dorsal root ganglia. *Eur J Pharmacol* 613: 54–59.
- Déciga-Campos M, Ramírez-Marín PM, López-Muñoz FJ (2015). Synergistic antinociceptive interaction between palmitoylethanolamide and tramadol in the mouse formalin test. *Eur J Pharmacol* 765: 68–74.
- De Filippis D, Luongo L, Cipriano M, Palazzo E, Cinelli MP, De Novellis V *et al.* (2011). Palmitoylethanolamide reduces granuloma-induced hyperalgesia by modulation of mast cell activation in rats. *Mol Pain* 7: 3–3.
- de Guglielmo G, Kallupi M, Scuppa G, Stopponi S, Demopoulos G, Gaitanaris G *et al.* (2014). Analgesic tolerance to morphine is regulated by PPAR γ . *Br J Pharmacol* 171: 5407–5416.
- De Novellis V, Luongo L, Guida F, Cristino L, Palazzo E, Russo R *et al.* (2012). Effects of intra-ventrolateral periaqueductal grey palmitoylethanolamide on thermoceptive threshold and rostral ventromedial medulla cell activity. *Eur J Pharmacol* 676: 41–50.
- Delerive P, De Bosscher K, Vanden Berghe W, Fruchart JC, Haegeman G, Staels B (2002). DNA binding-independent induction of I κ B α gene transcription by PPAR α . *Mol Endocrinol* 16: 1029–1039.
- Delerive P, Fruchart JC, Staels B (2001). Peroxisome proliferator-activated receptors in inflammation control. *J Endocrinol* 169: 453–459.
- Delerive P, Gervois P, Fruchart JC, Staels B (2000). Induction of I κ B α expression as a mechanism contributing to the anti-inflammatory activities of peroxisome proliferator-activated receptor- α activators. *J Biol Chem* 275: 36703–36707.
- Desvergne B, Wahli W (1999). Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev* 20: 649–688.
- Di Cesare Mannelli L, D'Agostino G, Pacini A, Russo R, Zanardelli M, Ghelardini C *et al.* (2013). Palmitoylethanolamide is a disease-modifying agent in peripheral neuropathy: pain relief and neuroprotection share a PPAR-alpha-mediated mechanism. *Mediators Inflamm* 2013: 328797.
- Di Cesare Mannelli L, Pacini A, Corti F, Boccella S, Luongo L, Esposito E *et al.* (2015). Antineuropathic profile of N-palmitoylethanolamine in a rat model of oxaliplatin-induced neurotoxicity. *PLoS ONE* 10: e0128080.
- Donvito G, Bettoni I, Comelli F, Colombo A, Costa B (2015). Palmitoylethanolamide relieves pain and preserves pancreatic islet cells in a murine model of diabetes. *CNS Neurol Disord Drug Targets* 14: 452–462.
- Donvito G, Wilkerson JL, Damaj MI, Lichtman AH (2016). Palmitoylethanolamide reverses paclitaxel-induced allodynia in mice. *J Pharmacol Exp Ther* 359: 310–318.
- Donvito G, Bagdas D, Toma W, Rahimpour E, Jackson A, Meade JA *et al.* (2017). The interaction between alpha 7 nicotinic acetylcholine receptor and nuclear peroxisome proliferator-activated receptor- α represents a new antinociceptive signaling pathway in mice. *Exp Neurol* 295: 194–201.
- Dreyer C, Krey G, Keller H, Givel F, Helftenbein G, Wahli W (1992). Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell* 68: 879–887.
- Floyd JS, Barbehenn E, Lurie P, Wolfe SM (2009). Case series of liver failure associated with rosiglitazone and pioglitazone. *Pharmacoepidemiol Drug Saf* 18: 1238–1243.
- Fu J, Gaetani S, Oveisi F, Lo Verme J, Serrano A, Rodriguez De Fonseca F *et al.* (2003). Oleylethanolamide regulates feeding and body weight through activation of the nuclear receptor PPAR-[alpha]. *Nature* 425: 90–93.
- Friedland SN, Leong A, Filion KB, Genest J, Lega IC, Mottillo S *et al.* (2012). The cardiovascular effects of peroxisome proliferator-activated receptor agonists. *Am J Med* 125: 126–133.
- Fuchs PN, Peng YB, Boyette-Davis JA, Uhelski ML (2014). The anterior cingulate cortex and pain processing. *Front Integr Neurosci* 8: 35.
- Gabrielsson L, Mattsson S, Fowler CJ (2016). Palmitoylethanolamide for the treatment of pain: pharmacokinetics, safety and efficacy. *Br J Clin Pharmacol* 82: 932–942.
- Garg S, Deshmukh Vishwajit R, Prasoon P (2017). Possible modulation of PPAR- γ cascade against depression caused by neuropathic pain in rats. *J Basic Clin Physiol Pharmacol* 28: 593–600.
- Ghavimi H, Charkhpour M, Ghasemi S, Mesgari M, Hamishehkar H, Hassanzadeh K *et al.* (2015). Pioglitazone prevents morphine antinociceptive tolerance via ameliorating neuroinflammation in rat cerebral cortex. *Pharmacol Rep* 67: 78–84.
- Ghavimi H, Hassanzadeh K, Maleki-Dizaji N, Azarfardian A, Ghasami S, Zolali E *et al.* (2014). Pioglitazone prevents morphine antinociception tolerance and withdrawal symptoms in rats. *Naunyn Schmiedebergs Arch Pharmacol* 387: 811–821.
- Gill N, Bijjem K, Sharma P (2013). Anti-inflammatory and anti-hyperalgesic effect of all-trans retinoic acid in carrageenan-induced paw edema in Wistar rats: involvement of peroxisome proliferator-activated receptor- β/δ receptors. *Indian J Pharmacol* 45: 278–282.
- Green S, Tugwood JD, Issemann I (1992). The molecular mechanism of peroxisome proliferator action: a model for species differences and mechanistic risk assessment. *Toxicol Lett*, 64–65 Spec No : 131–139.

- Griggs RB, Donahue RR, Adkins BG, Anderson KL, Thibault O, Taylor BK (2016). Pioglitazone inhibits the development of hyperalgesia and sensitization of spinal nociceptive neurons in type 2 diabetes. *J Pain* 17: 359–373.
- Griggs RB, Donahue RR, Morgenweck J, Grace PM, Sutton A, Watkins LR *et al.* (2015). Pioglitazone rapidly reduces neuropathic pain through astrocyte and nongenomic PPAR γ mechanisms. *Pain* 156: 469–482.
- Guida F, Luongo L, Marmo F, Romano R, Iannotta M, Napolitano F *et al.* (2015). Palmitoylethanolamide reduces pain-related behaviors and restores glutamatergic synapses homeostasis in the medial prefrontal cortex of neuropathic mice. *Mol Brain* 8: 47.
- Harding SD, Sharman JL, Faccenda E, Southan C, Pawson AJ, Ireland S *et al.* (2018). The IUPHAR/BPS Guide to PHARMACOLOGY in 2018: updates and expansion to encompass the new guide to IMMUNOPHARMACOLOGY. *Nucl Acids Res* 46: D1091–D1106.
- Hasegawa-Moriyama M, Kurimoto T, Nakama M, Godai K, Kojima M, Kuwaki T *et al.* (2013). Peroxisome proliferator-activated receptor-gamma agonist rosiglitazone attenuates inflammatory pain through the induction of heme oxygenase-1 in macrophages. *Pain* 154: 1402–1412.
- Hasegawa-Moriyama M, Ohnou T, Godai K, Kurimoto T, Nakama M, Kanmura Y (2012). Peroxisome proliferator-activated receptor-gamma agonist rosiglitazone attenuates postincisional pain by regulating macrophage polarization. *Biochem Biophys Res Commun* 426: 76–82.
- Hesslink JMK, Hekker TAM (2012). Therapeutic utility of palmitoylethanolamide in the treatment of neuropathic pain associated with various pathological conditions: a case series. *J Pain Res* 5: 437–442.
- Impellizzeri D, Di Paola R, Cordaro M, Gugliandolo E, Casili G, Morittu VM *et al.* (2016). Adelmidrol, a palmitoylethanolamide analogue, as a new pharmacological treatment for the management of acute and chronic inflammation. *Biochem Pharmacol* 119: 27–41.
- Issemann I, Green S (1990). Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347: 645–650.
- Iwai S, Maeda T, Kiguchi N, Kobayashi Y, Fukazawa Y, Ozaki M *et al.* (2008). Pioglitazone attenuates tactile allodynia and microglial activation in mice with peripheral nerve injury. *Drug Discov Ther* 2: 353–356.
- Jain V, Jaggi AS, Singh N (2009). Ameliorative potential of rosiglitazone in tibial and sural nerve transection-induced painful neuropathy in rats. *Pharmacol Res* 59: 385–392.
- Jennings EM, Okine BN, Roche M, Finn DP (2014). Stress-induced hyperalgesia. *Prog Neurobiol* 121: 1–18.
- Jhaveri MD, Richardson D, Robinson I, Garle MJ, Patel A, Sun Y *et al.* (2008). Inhibition of fatty acid amide hydrolase and cyclooxygenase-2 increases levels of endocannabinoid related molecules and produces analgesia via peroxisome proliferator-activated receptor-alpha in a model of inflammatory pain. *Neuropharmacology* 55: 85–93.
- Jia HB, Wang XM, Qiu LL, Liu XY, Shen JC, Ji Q *et al.* (2013). Spinal neuroimmune activation inhibited by repeated administration of pioglitazone in rats after L5 spinal nerve transection. *Neurosci Lett* 543: 130–135.
- Jia H, Xu S, Liu Q, Liu J, Xu J, Li W *et al.* (2016). Effect of pioglitazone on neuropathic pain and spinal expression of TLR-4 and cytokines. *Exp Ther Med* 12: 2644–2650.
- Jia H, Zhu S, Ji Q, Hui K, Duan M, Xu J *et al.* (2010). Repeated administration of pioglitazone attenuates development of hyperalgesia in a rat model of neuropathic pain. *Exp Clin Psychopharmacol* 18: 359–365.
- Loke YK, Kwok CS, Singh S (2011). Comparative cardiovascular effects of thiazolidinediones: systematic review and meta-analysis of observational studies. *BMJ* 342: d1309.
- LoVerme J, La Rana G, Russo R, Calignano A, Piomelli D (2005). The search for the palmitoylethanolamide receptor. *Life Sci* 77: 1685–1698.
- LoVerme J, Russo R, La Rana G, Fu J, Farthing J, Mattace-Raso G *et al.* (2006). Rapid broad-spectrum analgesia through activation of peroxisome proliferator-activated receptor- α . *J Pharmacol Exp Ther* 319: 1051–1061.
- Lyons DN, Zhang L, Danaher RJ, Miller CS, Westlund KN (2017). PPAR γ agonists attenuate trigeminal neuropathic pain. *Clin J Pain* 33: 1071–1080.
- Lyons DN, Zhang L, Pandya JD, Danaher RJ, Ma F, Miller CS *et al.* (2018). Combination drug therapy of pioglitazone and D-cycloserine attenuates chronic orofacial neuropathic pain and anxiety by improving mitochondrial function following trigeminal nerve injury. *Clin J Pain* 34: 168–177.
- Maeda T, Kiguchi N, Kobayashi Y, Ozaki M, Kishioka S (2008). Pioglitazone attenuates tactile allodynia and thermal hyperalgesia in mice subjected to peripheral nerve injury. *J Pharmacol Sci* 108: 341–347.
- Mansouri MT, Naghizadeh B, Ghorbanzadeh B, Alboghobeish S (2017a). Systemic and local anti-nociceptive effects of simvastatin in the rat formalin assay: role of peroxisome proliferator-activated receptor γ and nitric oxide. *J Neurosci Res* 95: 1776–1785.
- Mansouri MT, Naghizadeh B, Ghorbanzadeh B, Rajabi H, Pashmforoush M (2017b). Pharmacological evidence for systemic and peripheral antinociceptive activities of pioglitazone in the rat formalin test: role of PPAR γ and nitric oxide. *Eur J Pharmacol* 805: 84–92.
- Moreira DRM, Santos DS, Espirito Santo RFD, Santos FED, de Oliveira Filho GB, Leite ACL *et al.* (2017). Structural improvement of new thiazolidinones compounds with antinociceptive activity in experimental chemotherapy-induced painful neuropathy. *Chem Biol Drug Des* 90: 297–307.
- Moreno S, Farioli-Vecchioli S, Ceru MP (2004). Immunolocalization of peroxisome proliferator-activated receptors and retinoid X receptors in the adult rat CNS. *Neuroscience* 123: 131–145.
- Morgenweck J, Abdel-Aleem OS, Mcnamara KC, Donahue RR, Badr MZ, Taylor BK (2009). Activation of peroxisome proliferator-activated receptor gamma in brain inhibits inflammatory pain, dorsal horn expression of Fos, and local edema. *Neuropharmacology* 58: 337–345.
- Morgenweck J, Griggs RB, Donahue RR, Zadina JE, Taylor BK (2013). PPAR γ activation blocks development and reduces established neuropathic pain in rats. *Neuropharmacology* 70: 236–246.
- Murad H, Ayuob N (2015). Co-administration of pioglitazone improves fluoxetine's antinociceptive, neuroprotective, and antidepressant effects in chronic constriction injury in rats. *Pain Physician* 18: 609–620.
- Napimoga MH, Souza GR, Cunha TM, Ferrari LF, Clemente-Napimoga JT, Parada CA *et al.* (2007). 15d-Prostaglandin J₂ inhibits inflammatory hypernociception: involvement of peripheral opioid receptor. *J Pharmacol Exp Ther* 324: 313–321.

- Neugebauer V (2015). 15. Amygdala pain mechanisms. *Handb Exp Pharmacol* 227: 261–284.
- Nissen SE, Wolski K (2007). Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes. *New England Journal of Medicine* 356: 2457–2471.
- O'Mahony SM, Clarke G, Mckernan DP, Bravo JA, Dinan TG, Cryan JF (2013). Differential visceral nociceptive, behavioural and neurochemical responses to an immune challenge in the stress-sensitive Wistar Kyoto rat strain. *Behav Brain Res* 253: 310–317.
- Okine BN, Madasu MK, MCGowan F, Prendergast C, Gaspar JC, Harhen B *et al.* (2016). N-palmitoylethanolamide in the anterior cingulate cortex attenuates inflammatory pain behaviour indirectly via a CB1 receptor-mediated mechanism. *Pain* 157: 2687–2696.
- Okine BN, Gaspar JC, Madasu MK, Olango WM, Harhen B, Roche M *et al.* (2017). Characterisation of peroxisome proliferator-activated receptor signalling in the midbrain periaqueductal grey of rats genetically prone to heightened stress, negative affect and hyperalgesia. *Brain Res* 1657: 185–192.
- Okine BN, Rea K, Olango WM, Price J, Herdman S, Madasu MK *et al.* (2014). A role for PPAR α in the medial prefrontal cortex in formalin-evoked nociceptive responding in rats. *Br J Pharmacol* 171: 1462–1471.
- Okine BN, Spicer C, Millns P, Bennett A, Chapman V (2015). Systemic administration of WY-14643, a selective synthetic agonist of peroxisome proliferator activator receptor-alpha, alters spinal neuronal firing in a rodent model of neuropathic pain. *Scand J Pain* 9: 42–48.
- Olango WM, Finn DP (2014). Neurobiology of stress-induced hyperalgesia. *Curr Top Behav Neurosci* 20: 251–280.
- Oliveira AC, Bertollo CM, Rocha LT, Nascimento EB Jr, Costa KA, Coelho MM (2007). Antinociceptive and antiedematogenic activities of fenofibrate, an agonist of PPAR α , and pioglitazone, an agonist of PPAR γ . *Eur J Pharmacol* 561: 194–201.
- O'Sullivan SE, Kendall DA (2010). Cannabinoid activation of peroxisome proliferator-activated receptors: potential for modulation of inflammatory disease. *Immunobiology* 215: 611–616.
- Paragomi P, Rahimian R, Kazemi MH, Gharedaghi MH, Khalifeh-Soltani A, Azary S *et al.* (2014). Antinociceptive and antidiarrheal effects of pioglitazone in a rat model of diarrhoea-predominant irritable bowel syndrome: role of nitric oxide. *Clin Exp Pharmacol Physiol* 41: 118–126.
- Park SW, Yi JH, Miranpuri G, Satriotomo I, Bowen K, Resnick DK *et al.* (2007). Thiazolidinedione class of peroxisome proliferator-activated receptor gamma agonists prevents neuronal damage, motor dysfunction, myelin loss, neuropathic pain, and inflammation after spinal cord injury in adult rats. *J Pharmacol Exp Ther* 320: 1002–1012.
- Pena-dos-Santos DR, Severino FP, Pereira SAL, Rodrigues DBR, Cunha FQ, Vieira SM *et al.* (2009). Activation of peripheral κ/δ opioid receptors mediates 15-deoxy- Δ 12,14-prostaglandin J₂ induced-antinociception in rat temporomandibular joint. *Neuroscience* 163: 1211–1219.
- Pottabathini R, Kumar A, Bhatnagar A, Garg S, Ekavali E (2016). Ameliorative potential of pioglitazone and ceftriaxone alone and in combination in rat model of neuropathic pain: targeting PPAR γ and GLT-1 pathways. *Pharmacol Rep* 68: 85–94.
- Rea K, Olango WM, Okine BN, Madasu MK, Mcguire IC, Coyle K *et al.* (2014). Impaired endocannabinoid signalling in the rostral ventromedial medulla underpins genotype-dependent hyperresponsivity to noxious stimuli. *Pain* 155: 69–79.
- Ruiz-Medina, J., Flores, J.A., Tasset, I., Tunez, I., Valverde, O., AND Fernandez-Espejo, E. (2012). Alteration of neuropathic and visceral pain in female C57BL/6J mice lacking the PPAR- α gene. *Psychopharmacology (Berl)* 222: 477–488.
- Russo R, Caro CD, Avagliano C, Cristiano C, La Rana G, Mattace Raso G *et al.* (2016). Sodium butyrate and its synthetic amide derivative modulate nociceptive behaviors in mice. *Pharmacol Res* 103: 279–291.
- Russo R, LoVerme J, La Rana G, D'Agostino G, Sasso O, Calignano A *et al.* (2007). Synergistic antinociception by the cannabinoid receptor agonist anandamide and the PPAR-alpha receptor agonist GW7647. *Eur J Pharmacol* 566: 117–119.
- Sagar DR, Kendall DA, Chapman V (2008). Inhibition of fatty acid amide hydrolase produces PPAR-alpha-mediated analgesia in a rat model of inflammatory pain. *Br J Pharmacol* 155: 1297–1306.
- Saito T, Hasegawa-Moriyama M, Kurimoto T, Yamada T, Inada E, Kanmura Y (2016). Resolution of inflammation by resolvin D1 is essential for peroxisome proliferator-activated receptor- γ -mediated analgesia during postincisional pain development in type 2 diabetes. *Anesthesiology* 123: 1420–1434.
- Sasso O, Russo R, Vitiello S, Raso GM, D'Agostino G, Iacono A *et al.* (2012). Implication of allopregnanolone in the antinociceptive effect of N-palmitoylethanolamide in acute or persistent pain. *Pain* 153: 33–41.
- Smith AF, Macfie WG, Oliver MF (1970). Clofibrate, serum enzymes, and muscle pain. *Br Med J* 2: 86–88.
- Staels B, Koenig W, Habib A, Merval R, Lebret M, Torra IP *et al.* (1998). Activation of human aortic smooth-muscle cells is inhibited by PPAR α but not by PPAR γ activators. *Nature* 393: 790–793.
- Suardi az M, Estivill-Torr us G, Goicoechea C, Bilbao A, Rodr iguez De Fonseca F (2007). Analgesic properties of oleoylethanolamide (OEA) in visceral and inflammatory pain. *Pain* 133: 99–110.
- Sunder Mudaliar MF, Chang AR, Henry RR (2003). Thiazolidinediones, peripheral edema, and type 2 diabetes: incidence, pathophysiology, and clinical implications. *Endocrine Practice* 9: 406–416.
- Takahashi Y, Hasegawa-Moriyama M, Sakurai T, Inada E (2011). The macrophage-mediated effects of the peroxisome proliferator-activated receptor-gamma agonist rosiglitazone attenuate tactile allodynia in the early phase of neuropathic pain development. *Anesth Analg* 113: 398–404.
- Taylor BK, Dadia N, Yang CB, Krishnan S, Badr M (2002). Peroxisome proliferator-activated receptor agonists inhibit inflammatory edema and hyperalgesia. *Inflammation* 26: 121–127.
- Tugwood JD, Issemann I, Anderson RG, Bundell KR, Mcpheat WL, Green S (1992). The mouse peroxisome proliferator activated receptor recognizes a response element in the 5' flanking sequence of the rat acyl CoA oxidase gene. *EMBO J* 11: 433–439.
- Tuo W, Leleu-Chavain N, Spencer J, Sansook S, Millet R, Chavatte P (2017). Therapeutic potential of fatty acid amide hydrolase, monoacylglycerol lipase, and N-acylethanolamine acid amidase inhibitors. *J Med Chem* 60: 4–46.
- van Eekeren ICM, Clockaerts S, Bastiaansen-Jenniskens YM, Lubberts E, Verhaar JAN, Van Osch GJVM *et al.* (2013). Fibrates as therapy for osteoarthritis and rheumatoid arthritis? A systematic review. *Ther Adv Musculoskelet Dis* 5: 33–44.
- Vasconcelos MAL, Royo VA, Ferreira DS, Crotti AEM, e Silva MLA, Carvalho JCT *et al.* (2006). In vivo analgesic and anti-

inflammatory activities of ursolic acid and oleanoic acid from *Miconia albicans* (melastomataceae). *Z Naturforsch C* 61:477–482.

Wang J, Zhang Q, Zhao L, Li D, Fu Z, Liang L (2014). Down-regulation of PPAR α in the spinal cord contributes to augmented peripheral inflammation and inflammatory hyperalgesia in diet-induced obese rats. *Neuroscience* 278: 165–178.

Warden A, Truitt J, Merriman M, Ponomareva O, Jameson K, Ferguson LB *et al.* (2016). Localization of PPAR isotypes in the adult mouse and human brain. *Sci Rep* 6: 27618.

Wright MB, Bortolini M, Tadayyon M, Bopst M (2014). Minireview: challenges and opportunities in development of PPAR agonists. *Mol Endocrinol* 28: 1756–1768.

Yang L, Li L, Chen L, Li Y, Chen H, Li Y *et al.* (2015). Potential analgesic effects of a novel N-acyl ethanolamine acid amidase inhibitor F96 through PPAR- α . *Sci Rep* 5: 13565.

Zanardelli M, Micheli L, Cinci L, Failli P, Ghelardini C, Di Cesare Mannelli L (2014). Oxaliplatin neurotoxicity involves peroxisome alterations. PPAR γ agonism as preventive pharmacological approach. *PLoS One* 9: e102758.

Zhang, Y., Song, C., Li, H., Hou, J., AND Li, D. (2016). Ursolic acid prevents augmented peripheral inflammation and inflammatory hyperalgesia in high-fat diet-induced obese rats by restoring downregulated spinal PPAR α . *Mol Med Rep* 13: 5309–5316.