



Provided by the author(s) and University of Galway in accordance with publisher policies. Please cite the published version when available.

Title	The effect of operator position on the quality of chest compressions delivered in a simulated ambulance
Author(s)	Mullin, Scott; Lydon, Sinéad; O'Connor, Paul
Publication Date	2019-12-09
Publication Information	Mullin, Scott, Lydon, Sinéad, & O'Connor, Paul. (2020). The Effect of Operator Position on the Quality of Chest Compressions Delivered in a Simulated Ambulance. <i>Prehospital and Disaster Medicine</i> , 35(1), 55-60. doi:10.1017/S1049023X19005193
Publisher	Cambridge University Press
Link to publisher's version	https://doi.org/10.1017/S1049023X19005193
Item record	http://hdl.handle.net/10379/16296
DOI	http://dx.doi.org/10.1017/S1049023X19005193

Downloaded 2024-04-26T14:59:49Z

Some rights reserved. For more information, please see the item record link above.



The Effect of Operator Position on the Quality of Chest Compressions Delivered in a Simulated Ambulance

ABSTRACT

Background. Ambulances are where patient care is often initiated or maintained, but this setting poses safety risks for paramedics. Paramedics have been found that in order to optimize patient care, they must compromise their own safety by standing unsecured in a moving ambulance.

Hypothesis/Problem. To compare the quality of chest compressions (CCs) in the two positions they can be delivered in an ambulance.

Methods. A randomized counterbalanced study was carried out with 24 paramedic students. Simulated CCs were performed in a stationary ambulance on a cardiopulmonary resuscitation manikin for 2-minutes from either: (A) an unsecured standing position or (B) a seated secured position. Participant's attitudes toward the effectiveness of the two positions were evaluated.

Results. The mean total number of CCs were not significantly different standing unsecured (220; SD=12) as compared to seated and secured (224; SD =21). There was no significant difference in mean compression rate standing unsecured (110 compressions per minute; SD=6) as compared to seated and secured (113 compressions per minute; SD=10). CCs performed in the unsecured standing position yielded a significantly greater mean depth (52mm; SD= 6) than did seated secured (26mm; SD= 7; $p<.001$). Additionally, the standing unsecured position produced a significantly higher percentage (83%; SD= 21) for the number of correct compressions, as compared to the seated secured position (8%; SD= 17; $p<.001$). Participants also believed that CCs delivered from standing were more effective than those delivered when seated.

Conclusions. The quality of CCs delivered from a seated and secured position is inferior to those delivered from an unsecured standing position. There is a need to consider how training, technologies, and ambulance design can impact the quality of CCs.

Keywords: Cardiopulmonary resuscitation; simulation; Ambulance; chest compressions; Out-of-hospital cardiac arrest.

The Effect of Operator Position on the Quality of Chest Compressions Delivered in a Simulated Ambulance

INTRODUCTION

Ambulances are clinical environments in which patient care is initiated or maintained, but this setting poses safety risks for paramedics during transport. It has been suggested that there is a 'hidden crisis' in the emergency medical industry, with U.S paramedics 2.5 times more likely to die on the job as compared to the U.S. national average.¹ A twenty-year review by the United States National Highway Traffic Safety Administration (NHTSA) identified that, on average, 4,500 ambulances are involved in crashes annually, with 34% resulting in injuries to paramedics.² Moreover, 84% of paramedics in the crashes were unrestrained within the patient compartment of the ambulance.² Emergency Medical Services (EMS) have seatbelt policies for the patient compartment, with paramedics mandated to wear seatbelts at all times.³ However, it is standard policy to permit the removal of any restraint when the restraint either delays or prevents clinical care.³

Cardio pulmonary resuscitation (CPR) is one key example of a clinical intervention frequently performed by paramedics during ambulance transport in which the use of a seatbelt may negatively impact patient care. Effective chest compressions are a key feature of high-quality CPR,⁴⁻⁶ and in an ambulance may be delivered in either a standing or sitting position. Effective chest compressions are characterised by compressions to the correct depth, rate, and full recoil.⁷ However, despite the importance of delivering effective chest compressions, there are relatively few studies that have examined the impact of the body position of the individual delivering the CPR on the quality of chest compressions.

Paramedics are trained to give chest compressions as specified by the BLS guidelines within

standardized CPR courses, from a kneeling position on the floor. Although this position is standardized and meets current guidelines for effective compressions, there is no formal standard to address the delivery of compressions in a moving ambulance.

In pre-hospital simulated environments, no significant difference has been found in the quality of chest compressions delivered based on an operator that is either standing or kneeling.^{8,9} However, there is a dearth of research that directly compares the quality of chest compressions delivered in an ambulance setting between a standing and unsecured and a seated and secured position. This data is important for considering how to deliver optimum care to patients in an ambulance while preserving the safety of the paramedic.

The effectiveness of chest compressions from two conditions will be compared during simulated resuscitation in a stationary ambulance: from a standing unsecured position, or from a seated and secured position. The participants' preference for body position when delivering chest compressions in an ambulance will also be assessed. The implications for paramedic training, ambulance design, and the use of technologies such as mechanical CPR devices and real-time CPR feedback will be discussed.

METHODS


This study was conducted and reported in accordance with the reporting guidelines for health care simulation research reporting.¹⁰

Design

The study utilized a randomized counterbalanced design. The conditions for this study consisted of (A) standing unsecured position, and (B) seated and secured position (see Figure

1). In condition A, the paramedic is standing with knees slightly bent with straight arms on the chest of the patient, and uses their body weight to help deliver the compressions. In condition B, the paramedic is seated and must lean out over the patient in order to deliver the compressions.

Figure 1 Condition A- standing unsecured position, and Condition B- seated and secured position.



Setting

The study was carried out at the simulation centre of a Canadian college. The population of participants was 60 students across the first and second-year cohorts of the college's paramedic programme. Participants were excluded if they were not first or second-year students in the college's paramedic programme.

First year students have approximately 8 hours of experience performing CPR on high and low fidelity mannequins, and second year students have approximately 20 hours of experience. To be eligible for inclusion, the participants were required to be a student on the

programme and to hold a current certificate in Basic Life Support (BLS). The data collection was carried out between January and March 2018.

Ethical Approval

This study was reviewed and approved by the Medicine Hat College, Canada, Ethics Review Board.

Sample size calculation

No previous published papers were available specifically comparing the quality of chest compressions given from seated and standing positions. Therefore, the effect size from Sebbane et al¹¹ which compared kneeling and standing chest compressions was used to estimate the sample size required to adequately power the current study using GPower. Specifying an effect size of Cohen's $d=0.8$, a two-tailed matched-pairs t-test, alpha of .05, and power of 0.95 it was determined that it was necessary to recruit a minimum of 23 participants in order for the study to be sufficiently powered.

Equipment

Chest compressions were performed on a QCPR manikin (Laerdal, Canada) placed on an MX-PRO R3 Stryker ambulance cot (Stryker EMS, USA). The manikin has been used in a number of other studies of the quality of chest compressions.^{7,11,12} The Stryker stretcher was used as it is the current cot used within in most ambulances in Alberta, Canada. The simulation manikin and ambulance cot were placed and secured within a stationary simulated ambulance patient compartment (SimLeader, Canada). This simulation compartment is consistent with the specifications and functionality of the Alberta ambulance currently in service.

Measures

Chest compression data was captured for each participant on a Laerdal SimPad Plus SkillReporter (Laerdal, Canada). The participants performed compressions on a Resusci Anne Q CPR manikin (Laerdal, Canada). All data was downloaded onto a laptop and saved into the Session Viewer software provided by Laerdal. Each participant was assigned a random code to blind the research team as to which condition the participant had completed first. The data collected were: the total number of compressions; the number of correct compressions (compressions performed at the correct rate, depth, with hands in the correct position and with full release of pressure at the end of compression); and mean compression depth, and average rate.

Preference and perception data by participants was assessed by using a 5-point Likert scale from 1 ('strongly disagree') to 5 ('strongly agree') via an online survey in Google Forms.

Participants responded to the following statements:

- the chest compressions I performed when seated and secured were effective;
- the chest compressions I performed when standing unsecured were effective;
- I prefer the seated and secured body position when performing chest compressions;
and
- I prefer the standing unsecured body position when performing chest compressions.


Procedure

Students in the first and second-year cohorts in a paramedic programme at a Canadian College were invited to participate in January 2018. The students received written information about the study via email, and were given the opportunity to ask questions. The

first 24 students who agreed to participate were recruited into the study. All participants provided written informed consent.

The participant flow diagram in Figure 2 provides an overview of the stages of the study. Participants were randomized into the two groups by using an online random number generator (www.graphpad.com). The research team were blinded to the randomisation. As the research team only had access to one ambulance simulator, the two groups started at alternative times on the same day. All participants completed a 30-minute refresher activity to ensure adequate knowledge of the Canadian Heart and Stroke Foundation BLS CPR before data collection. Each participant subsequently performed a two-minute chest compression routine on the Resusci Anne Quality Cardiopulmonary Resuscitation (Q CPR) manikin, with visual feedback, to practice prior to the data collection.

Figure 2. Participant flow diagram.



Following orientation, individual participants entered the ambulance simulator and were asked to perform continuous chest compressions for two minutes followed by a 10-minute

break between conditions. Participants were in a standing unsecured position beside an ambulance cot/stretchers (condition A; see Figure 1) and alternatively seated in the seated and secured position (condition B; see Figure 1). Participants did not receive visual or audio feedback on the quality of the chest compressions in either condition.

Following completion of testing in both conditions, participants were asked to complete an anonymous online survey indicating their agreement to four questions relating to their preference for delivering chest compressions outlined in the methods section.

Statistical Analysis

IBM SPSS (version 23) was used to complete the statistical analyses. Comparisons between the two CPR positions were performed by using the dependent t-test or Wilcoxon Signed Rank test when data was not normally distributed. Comparisons were made across the following dimensions: percent of correct compressions, the average rate of compressions per minute, the average depth of compressions, and number of total compressions. A Bonferroni correction was applied to control for the impact of multiple comparisons and a significance level of .012 was adopted. Effect size was calculated using Cohen's *d*.

RESULTS

Participants

Twenty four participants were recruited (11 first year and 13 second year students). Twelve participants were randomised to condition A, and 12 participants to condition B (see Figure 2). Fourteen participants were male and 10 were female.

Evaluation of operator position on chest compressions

There was no significant difference in the total number of compressions, or in the mean rate of compressions between the standing and seated positions (see Table 1). However, mean compression depth was significantly higher in the standing position as compared to the seated position, with a large effect size of the difference (see Table 1). There was also a significant difference, and large effect size of the difference, in the percentage of correct compressions between the two positions (see Table 1). The standing position resulted in a significantly higher percentage of correct compressions (total number of compressions, compressions performed at the correct rate, depth, with hands in the correct position and with full release of pressure at the end of compression) than the seated position (see Table 1).

Table 1. Chest compression parameters according to operator position

Position	Standing (n =24)		Seated (n =24)		Significance (<i>p</i>)	Effect size (<i>d</i>)
	Mean	St dev	Mean	St dev		
Total chest compressions	220	12	224	21	n.s.* (0.36)	0.23
Rate (n/min)	110	6	113	10	n.s. (0.16)	0.36
Depth (mm)	52	6	26	7	<.001	3.98
Correct Compressions (%)	83	21	8	17	<.001	3.93

Attitudes to operator position

The participants perceived that the effectiveness of the chest compressions they delivered was higher when standing as compared to seated, and preferred to give chest compressions from a standing position (see Table 2).

Table 2. Likert-Scale Preference and Perception Questions (1 ‘strong disagree’ to 5 ‘strongly agree’; n=24).

Question	Median	Interquartile range	
		Lower (Q ₁)	Upper (Q ₃)
The chest compressions I performed when seated and secured were effective.	1	1	1
The chest compressions I performed when standing unsecured were effective.	5	4	5
I prefer the seated and secured body position when performing chest compressions.	1	1	1
I prefer the standing unsecured body position when performing chest compressions.	5	5	5

DISCUSSION

Delivering effective chest compressions during a cardiopulmonary arrest is crucial.⁴⁻⁶ However, this is a task that can be challenging to both effectively and safely perform in an ambulance. The study reported in this paper is one of a few studies that have specifically examined the impact of body position of the paramedic on the quality of chest compressions- and the only study of which the authors are aware that compares standing versus seated positions. The data indicate that delivering chest compressions from a seated and secured position, although safer for the paramedic, compromises the quality of the chest compressions given to the patient. Therefore, to optimize patient care, the paramedic must compromise their own safety by standing unsecured in the moving ambulance. These findings have implications for the training of paramedics to deliver chest compressions, ambulance design, and the use of technologies such as mechanical CPR devices, and real-time feedback.

It is encouraging to see that, when standing, the mean depth of the compressions given by the participants in this study exceeded the AHA guidelines of 50mm,¹³ and the percentage of correct compressions was greater than 80%. Moreover, the total number of compressions and the mean rate of compressions were not significantly different between the standing and seated positions. Other studies have also found that rate of compression is not effected by the position from which the chest compressions are delivered.^{7,11,14,15} This finding suggests that the chest compression rate learned during practice generalizes to different delivery positions. However, there would appear to be a trade-off in in which the depth of the compression is compromised in order for the paramedic to achieve an effective rate in all positions.

Chest compressions delivered from a seated position were significantly shallower, and had a significantly lower percentage of correct compressions, as compared to chest compressions delivered from a standing position. The participants' seated and secured mean compression depth of 26mm was comparable to the 27mm depth reported by Sebbane et al¹¹ in the evaluation of the effect of delivering kneeling chest compressions to a patient on a stretcher. However, in both cases the depth is considerably less than the AHA guidelines.¹³ Similarly, the percentage of correct compressions of less than 10% in the seated position is of great concern.

It is possible to speculate that this low number of correct compressions can be attributed to a lack of training and practice in delivering chest compressions from a seated position. A study that compared CPR training given with a mannequin on the floor, as compared to in a bed, found that the depth of compressions was too low in both cases. It was concluded that participants who optimized their working position (e.g. by lowering the height of the bed) performed deeper chest compressions)¹⁵. However, there are limited options in an ambulance

to change the environment in order to improve the quality of the chest compression other than standing in an unsecured position. Moreover, a bigger issue than training and practice is that the delivery of effective chest compressions from a seated position has a number of biomechanical challenges. When standing in an ambulance chest compressions are carried out through movement of the elbow and shoulder joints, and when kneeling through flexion and extension of the hip joint.⁸ When seated, all of the movement will have to come from the shoulder joints. Therefore this will likely lead to provider fatigue.

Even in the absence of any feedback, the participants were aware that chest compressions delivered while standing were more effective than when seated. Therefore, unsurprisingly, the participants preferred to give chest compressions from a standing position. Although this can only be inferred, these attitudes suggest that the participants are likely to deliver chest compressions from a standing position in an ambulance. A qualitative study carried out with Swedish paramedics found that they have an attitude of ‘patient first, safety second’ and prioritise the care of critically ill patients over their own safety in a moving ambulance.¹⁶ This suggests that paramedics are willing to compromise their own safety in order to deliver more effective patients care.

The findings from the study reported in this paper have a number of implications for future research directions. Devices that provide real-time feedback on the quality of chest compressions are becoming more commonplace. Feedback can be given by a metronome, voice prompts, visual displays, or digital timers.¹⁷ It has been found that verbal and visual feedback improves CPR quality in flying helicopters and moving ambulance vehicles.⁹ However, even with training, practice, and feedback we believe it is unlikely that it is possible to deliver good quality chest compressions from a seated and secured position.

Therefore, future research should consider other solutions to the safe delivery of chest compressions in an ambulance.

When transporting patients, where high-quality chest compressions cannot be delivered safely, the use of mechanical CPR may be a potential solution.¹⁸ Mechanical chest compression devices are designed to perform chest compressions at a specified rate and depth. Current research suggests that there is no significant difference in patient outcomes between mechanical and manual compressions when used in the pre-hospital environment.¹⁸ Therefore, assessing the use of mechanical CPR devices in ambulance transportation is warranted. However, this research should not only consider the outcome for patients, but also the impact of the use of mechanical CPR devices on the safety of the paramedic.

Research on CPR in hospital environments has found that providers can improve quality of chest compressions by making changes to the environment such as lowering the bed.¹⁵ Current configurations of ambulances provide limited opportunity to make changes to optimise the delivery of chest compressions. However, consideration should be given to changes that could be made (e.g. the ability to raise the seat) in order to allow the paramedic to optimising their position for delivering chest compressions. A greater focus on paramedic safety during transport is long overdue. Understanding how paramedics work within an ambulance may address inefficiencies in design and functionality that contribute to the potential for injury to the paramedic and compromise patient care.¹⁹

This study has a number of limitations. First, the participants were paramedic students rather than experienced and qualified paramedics. Nevertheless, from an experimental perspective, a student group is desirable as all participants have a similar level of experience and training. Second, the participants performed chest compressions for two minutes, which is short and

not representative of a full resuscitation. It is known that the quality of chest compressions decrease over time.²⁰ However, the data reported in this paper does represent one cycle of CPR, and the time is consistent with other studies that have examined the impact of operator position on quality of chest compressions.¹¹ Third, the number of participants in the study was relatively low. However, in spite of the low number of participants, the study was adequately powered, and the quantity of participants was larger than other similar studies.⁷⁻⁹ Finally, the study used a manikin and a simulated stationary ambulance. Participants were not challenged by acceleration, deceleration, and G-forces inherent with a moving ambulance.²¹

CONCLUSION

The quality of chest compressions delivered from a seated and secured position are inferior to those delivered from an unsecured standing position. This demonstrates that paramedics must compromise their own safety in order to deliver effective chest compressions in an ambulance. There is a need to consider how training, technologies, and the design of the patient compartment of ambulances can impact not only patient care, but also and paramedic safety in ambulance transfers. Paramedics should not be placed in a situation in which they must compromise their own safety for patient care.

Acknowledgements: We would like to acknowledge A. Woods (Medicine Hat College, Medicine Hat, AB) for drawing the two illustrations in Figure 1.

Disclosure statement: The authors report no conflict of interest.

Contributors: SW, SL, and POC designed the study. SW collected the data. SW, SL, and POC analysed and interpreted the data and drafted the manuscript. All authors approved the draft manuscript prior to submission.

Funding: The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Ethics approval: This study was reviewed and approved by the Medicine Hat College, Canada, Ethics Review Board.

REFERENCES

1. Maguire BJ, Hunting KL, Smith GS, Levick NR. Occupational fatalities in emergency medical services: a hidden crisis. *Ann Emerg Med* 2002;40(6):625-32.
2. Smith N. A National Perspective on Ambulance Crashes and Safety. Guidance from the National Highway Traffic Safety Administration on ambulance safety for patients and providers. *EMS World* 2015;44(9):91-2, 4.
3. Alberta Health Services EMS. Securing EMS staff, patients, passengers and equipment in ground vehicles: corporate Policy and procedure (PS-EMS-01). Alberta: Author, 2016.
4. Wik L, Kramer-Johansen J, Myklebust H, Sorebo H, Svensson L, Fellows B, et al. Quality of cardiopulmonary resuscitation during out-of-hospital cardiac arrest. *JAMA* 2005;293(3):299-304.
5. Sutton RM, Nadkarni V, Abella BS. "Putting it all together" to improve resuscitation quality. *Emerg Med Clin North Am* 2012;30(1):105-22.
6. Cheskes S, Byers A, Zhan C, Verbeek PR, Ko D, Drennan IR, et al. CPR quality during out-of-hospital cardiac arrest transport. *Resuscitation* 2017;114:34-9.
7. Perkins GD, Smith CM, Augre C, Allan M, Rogers H, Stephenson B, et al. Effects of a backboard, bed height, and operator position on compression depth during simulated resuscitation. *Intensive Care Med* 2006;32(10):1632-5.
8. Yasuda Y, Kato Y, Sugimoto K, Tanaka S, Tsunoda N, Kumagawa D, et al. Muscles used for chest compression under static and transportation conditions. *Prehosp Emerg Care* 2013;17(2):162-9.
9. Havel C, Schreiber W, Trimmel H, Malzer R, Haugk M, Richling N, et al. Quality of closed chest compression on a manikin in ambulance vehicles and flying helicopters with a real time automated feedback. *Resuscitation* 2010;81(1):59-64.

10. Cheng A, Kessler D, Mackinnon R, Chang TP, Nadkarni VM, Hunt EA, et al. Reporting guidelines for health care simulation research: extensions to the CONSORT and STROBE statements. *Advanc Sim* 2016;1(1):25.
11. Sebbane M, Hayter M, Romero J, Lefebvre S, Chabrot C, Mercier G, et al. Chest compressions performed by ED staff: a randomized cross-over simulation study on the floor and on a stretcher. *Am J Emerg Med* 2012;30(9):1928-34.
12. Cason CL, Trowbridge C, Baxley SM, Ricard MD. A counterbalanced cross-over study of the effects of visual, auditory and no feedback on performance measures in a simulated cardiopulmonary resuscitation. *BMC Nurs* 2011;10:15.
13. American Heart Association. Highlights of the 2015 American Heart Association guidelines update for CPR and ECC. Dallas, TX: Author, 2015.
14. Lewinsohn A, Sherren PB, Wijayatilake DS. The effects of bed height and time on the quality of chest compressions delivered during cardiopulmonary resuscitation: a randomised crossover simulation study. *Emerg Med J* 2012;29(8):660-3.
15. Mygind-Klausen T, Jaeger A, Hansen C, Aagaard R, Krogh LQ, Nebsbjerg MA, et al. In a bed or on the floor? - The effect of realistic hospital resuscitation training: A randomised controlled trial. *Am J Emerg Med* 2018;36(7):1236-41.
16. Suserud BO, Jonsson A, Johansson A, Petzall K. Caring for patients at high speed. *Emerg Nurs* 2013;21(7):14-8.
17. Brinkrolf P, Lukas R, Harding U, Thies S, Gerss J, Van Aken H, et al. A better understanding of ambulance personnel's attitude towards real-time resuscitation feedback. *Int J Qual Health Care* 2018;30(2):110-7.
18. Poole K, Couper K, Smyth MA, Yeung J, Perkins GD. Mechanical CPR: Who? When? How? *Crit Care* 2018;22(1):140.

19. Du B, Boileau M, Wierts K, Hignett S, Fischer S, Yazdani A. Existing Science on Human Factors and Ergonomics in the Design of Ambulances and EMS Equipment. *Prehosp Emerg Care* 2019;1-16.
20. Hightower D, Thomas SH, Stone CK, Dunn K, March JA. Decay in quality of closed-chest compressions over time. *Ann Emerg Med* 1995;26(3):300-3.
21. Slattery DE, Silver A. The hazards of providing care in emergency vehicles: an opportunity for reform. *Prehosp Emerg Care* 2009;13(3):388-97.