



Provided by the author(s) and NUI Galway in accordance with publisher policies. Please cite the published version when available.

| | |
|-----------------------------|---|
| Title | The impact of chemical amendment of dairy cattle slurry before land application on soil phosphorus dynamics |
| Author(s) | Brennan, Raymond B.; Healy, Mark G. |
| Publication Date | 2014-08-22 |
| Publication Information | Brennan, R.B., Wall, D., Fenton, O., Grant, J., Sharpley, A.N., Healy, M.G. (2014) 'The impact of chemical amendment of dairy cattle slurry before land application on soil phosphorus dynamics'. Communications In Soil Science And Plant Analysis, 45 (16):2215-2233. |
| Publisher | Taylor and Francis |
| Link to publisher's version | http://dx.doi.org/10.1080/00103624.2014.912293 |
| Item record | http://hdl.handle.net/10379/4515 |
| DOI | http://dx.doi.org/10.1080/00103624.2014.912293 |

Downloaded 2021-03-08T01:20:55Z

Some rights reserved. For more information, please see the item record link above.



Published as: Brennan, R.B., Wall, D., Fenton, O., Grant, J., Sharpley, A.N., Healy, M.G. 2014. The impact of chemical amendment of dairy cattle slurry before land application on soil phosphorus dynamics. Communications in Soil Science and Plant Analysis 45(16): 2215 – 2233.

DOI:10.1080/00103624.2014.912293

**Impact of chemical amendment of dairy cattle slurry on soil phosphorus dynamics
following application to soil**

R.B. Brennan^{a,b,c*}, D. Wall^b, O. Fenton^b, J. Grant^d, A.N. Sharpley^c, M.G. Healy^a

^aCivil Engineering, National University of Ireland, Galway, Co. Galway, Rep. of Ireland.

^bTeagasc, Environmental Research Centre, Johnstown Castle, Co. Wexford, Rep. of Ireland.

^cDept. Crop, Soil, and Environmental Sciences, Division of Agriculture, University of Arkansas,
Fayetteville, AR, USA.

^dTeagasc Research Centre, Ashtown, Co. Dublin, Rep. of Ireland.

*Corresponding author: Tel: +1 479 575 5720; 115 Plant Science Building, University of
Arkansas, Fayetteville, Arkansas, 72701; E-mail address: rbrennan@uark.edu

Abstract

A nine month incubation study was conducted to investigate the effectiveness of amending dairy cattle slurry with either alum, lime, poly-aluminum chloride (PAC), or ferric chloride (FeCl_3), in reducing water extractable P (WEP) levels in five soils (four mineral and one organic). Alum, lime, and PAC were the most effective amendments in decreasing WEP (compared to a slurry-control) for the four mineral soils (by an average of 47% at the end the 9-month incubation period). In comparison, FeCl_3 increased WEP (compared to the slurry-control) by an average of 35% at the end the study. None of the amendments examined effectively reduced WEP of the organic soil. No amendment reduced soil test P [(Morgan's P (P_m) and Mehlich-3 P (M3P)] compared to the soil-only treatment. Alum maintained the highest levels of M3P across the four mineral soils with the least risk of P loss to overlying water.

Keywords Water extractable phosphorus, soil test phosphorus, alum, lime, poly-aluminum chloride, management practices

Introduction

Land application of dairy cattle slurry can result in incidental and chronic phosphorus (P) losses to a waterbody (Buda et al. 2009), which may lead to eutrophication (Carpenter et al. 1998).

Incidental P losses occur when a rainfall event occurs shortly after slurry application and runoff is generated before the slurry has infiltrated the soil, while chronic P losses are long-term losses of P from soil as a result of a build-up in soil test P (STP) caused by the repeated or over-

application of inorganic fertilizers and manure (Preedy et al. 2001; Buda et al. 2009). The present study focuses on P loss from soils receiving chemically-amended dairy cattle slurry. When dairy cattle slurry is applied to soil, it results in an increase in STP (Sharpley et al. 2004). Once STP levels increase above agronomic optima, there is a higher risk of P loss to water bodies than from soils with lower STP levels (Sharpley et al. 1996; 2004). Water bodies near farms with a limited land base to distribute slurry are at particular risk, as slurry is repeatedly applied to high STP soils, are at particular risk (Doody et al. 2012; Wall et al. 2013). In these situations, both long- and short-term P management measures, such as implementing P drawdown strategies, transporting P off-farm, and soil, slurry and manure amendments that sequester P, may need to be utilized.

Chemical amendments have been shown to be effective in decreasing P solubility in dairy cattle and swine slurries (Dao and Daniel, 2002; Dou et al. 2003; Brennan et al. 2011a) and in mitigating incidental P losses in runoff (Smith et al. 2001; Elliot et al. 2005; Torbert et al. 2005; Brennan et al. 2011b; O' Flynn et al. 2012; Brennan et al. 2012). There has been limited research on the effect of chemical amendments used to lower soil P solubility on long-term P dynamics (Ann et al. 2000; Callahan et al. 2002; Kalbasi and Karthikeyan 2004; Moore and Edwards 2007). Kalbasi and Karthikeyan (2004) examined a silt loam soil with three different STP levels (12, 66, and 94 mg kg⁻¹ Bray-1 P, respectively) in an incubation experiment conducted over 24 months. An untreated slurry control and slurry amended with either alum, ferric chloride (FeCl₃), or lime were added to the three different STP soils. Kalbasi and Karthikeyan (2004) found that the effect of chemical amendment depended on treatment type, P application rate and background STP level, and recommended that more work was needed to investigate the

effectiveness of amendments in soils varying in physical and chemical characteristics. Callahan et al. (2002) amended four loam soils with seven P sorbing amendments and found that gypsum and water treatment residual (WTR) effectively decreased soil water extractable phosphorus (WEP) without any adverse agronomic effects. Moore and Edwards (2007) examined the effects of chemical amendment of poultry litter on a silt loam over a 20-year study duration. They found that long-term land application of alum-amended poultry litter decreased P runoff and that aluminum (Al) availability was lower from plots receiving alum-treated poultry manure than plots receiving ammonium ($\text{NH}_4\text{-N}$) fertilizer. Therefore, there is a need to examine how amendments perform across different soil types in order to identify situations (soil types, STP levels, climate, etc.) where they may be most effective.

Soil type plays a significant role in soil P solubility and it is critical that soil type is considered when examining the potential of amendments to reduce chronic P losses. Tunney (2000) found a strong association between STP (measured using Morgan's extracting solution) and dissolved reactive P (DRP) concentrations in overland flow in Irish grassland soils. This relationship can vary between different hydrological conditions (Kurz et al. 2005) and soil types (Daly et al. 2001; Regan et al. 2010). Daly et al. (2001) examined eleven soils chosen to best represent important agricultural grassland soils in Ireland varying in parent material, drainage, soil type and chemical properties, and found that, although STP was an important factor controlling P desorption, soil type also affected levels of sorption and desorption. Of particular interest is the effectiveness in organic soils, which are commonly found in parts of Ireland, mainland Europe and the USA. Such soils are often intensively farmed and require P applications annually, as they

have poor P retention capacity. Amendments may also offer the potential to apply slurry to peat soils with reduced risk of P losses and improved plant availability.

The objectives of this study were to investigate (1) the effectiveness of chemical amendments to reduce the solubility of P in dairy cattle slurry across five different Irish soils with different physical and chemical characteristics, (2) changes and relationships between soil WEP and STP following incorporation of amended slurry into soil under controlled laboratory conditions, and (3) to use the results to make recommendations on which soils, or soil properties, can be used to select locations within catchments most suitable for strategic use of chemical amendments.

Materials and Methods

Soil collection and analysis

Soils with different texture, organic matter (OM), and pH were selected to test the effectiveness of the amendments in a variety of conditions, and to represent some common soil types used in dairy farming in Ireland. The five soils collected were in the agronomic optimum STP range (Morgan's P levels of 5.1 to 8 mg L⁻¹) for productive grasslands with the exception of Soil C (Table 1), which was in the P-deficient range (< 5.0 mg L⁻¹) and the peaty soil (Soil E; 24.6 mg L⁻¹), which had an excessive STP level (>8.0 mg L⁻¹). The STP index system used in Ireland, ranges from 1 to 4, where a soil with P index 1 (very low) and 2 (low) has a likely response to P fertilizers, a P index 3 (agronomic optimum), and P index 4 (excessive) has no response to additional P fertilizer application and has a high risk of P loss. A peat soil was included, as there is a particular risk of P loss in runoff from peat soils due to their limited capacity to absorb P (Iyamuremye and Dick 1996; Cummins and Farrell 2003).

The soils were collected from five grassland sites, air dried, and crushed to pass a 2-mm sieve. Plant available soil P was determined by Morgan's P extract, the standard agronomic soil P testing method in Ireland (Morgan 1941) and by Mehlich-3 extractant (M3P) (Mehlich 1984). Mehlich-3 Al and iron (Fe) (M3-Al, and M3-Fe) were used to estimate degree of P saturation in the soils (Maguire and Sims 2002). Mehlich-3 calcium (Ca), cobalt (Co), copper (Cu), potassium (K), magnesium (Mg), manganese (Mn) and zinc (Zn) were also analysed. Soil WEP was determined after McDowell and Sharpley (2001) using an 100:1 deionised water-to-soil ratio.

Soil pH (n=3) was determined using a pH probe (WTW, Germany) and a 2:1 ratio of deionised water-to-soil. Particle size distribution (PSD) was determined using hydrometer method (B.S.1377-2:1990; British Standards 1990a) and the OM concentration of soil determined using loss of ignition (LOI) test (B.S.1377-3; British Standards 1990b). The results are presented in Table 1.

The approximate field capacity for each soil was determined after Bond et al. (2006). The approximate bulk density of each soil used in this study was determined based on the container volume occupied by 100 g of sieved soil after field capacity was achieved (Table 1).

Slurry collection and analysis

Slurry from dairy replacement heifers was taken from a farm (53°18' N, 8°47' W) in County Galway, Republic of Ireland, in June, 2010. The slurry in the storage tanks was fully agitated (mixed) prior to sampling and the slurry samples were transported to the laboratory in 10-L

drums. Slurry samples were stored for 48 h at 4°C prior to the incubation study and analysis was completed within 24 h of slurry being added to soil. Slurry pH was determined using a pH probe (WTW, Germany). The total P (TP), total N (TN) and total K (TK) of the dairy cattle slurry was determined following digestion of 3 g of well mixed slurry in a 100-mL Kjeldahl digestion flask with 10 mL of concentrated sulfuric acid (H₂SO₄) and 6 mL of 100 volumes hydrogen peroxide (H₂O₂) with selenium lithium sulphate tablets as a catalyst for 2 h before diluting the cooled digestate to 100 mL and filtering through No. 2 Whatman filter paper. Analyses for TP and TN were carried out colorimetrically using an automatic Multi-Channel Segmented Flow Analyzer (Burkard, United Kingdom) after Basson et al. (1968). Total K was analyzed using a Varian Spectra 400 Atomic Absorption (Varian, Switzerland). The WEP of slurry and amended slurry was measured at the time of soil application after Kleinman et al. (2007).

The slurry application was based on an application rate of 33 m³ ha⁻¹, which is common practice on grassland in Ireland (Coulter and Lalor 2008). This was equivalent to an application of 37.4 kg TP ha⁻¹ and approximately 0.94 kg WEP ha⁻¹. The same slurry application rate was selected for all soils to allow convenient randomisation. Amendments were applied at a stoichiometric ratio based on the TP of the slurry to the metal in the amendment. Alum was applied at 1.5:1 (Al:TP), lime at 16:1 (Ca:TP), poly-aluminium chloride (PAC) at 1.4:1 (Al:TP), and FeCl₃ at 1.5:1 (Fe:TP).

Incubation experiment

This 9-month incubation study comprised six treatments, conducted in triplicate, at a fixed temperature of 11°C on: (1) soil-only (to take account of the effects of incubation on soil P), (2)

soil mixed with dairy cattle slurry (slurry-control); and soil mixed with dairy cattle slurry, which was amended with either (3) alum, (4) lime, (5) PAC, or (6) FeCl_3 .

First, 100 g samples of air dried soil, passed through a 2-mm sieve, were placed in 0.5-L square containers (70 × 70 mm base). Then, slurry, or amended slurry, was added to soil and mixed before adding the volume of de-ionised water required to achieve 50% approximate field capacity (after Bond et al. 2006). The mineral soils had a mean bulk density of $1.12 \pm 0.15 \text{ g cm}^{-3}$ (Table 1). For the purpose of selecting slurry application rate, it was assumed that surface-applied slurry only interacts with the upper 20 mm of soil (Andraski et al. 2003; Sharpley 2003). Although the peat (Soil E) had a significantly different bulk density, slurry was applied at the same rate to the peat for consistency. Following this, the soil, slurry and water mixture was compacted using a custom-made packer to an appropriate bulk density similar to that in the field situation (Table 1). The containers were covered with para-film and perforated for air circulation. Throughout this study, the containers were weighed and water was added intermittently to ensure that approximately 50% field capacity was maintained during the incubation.

After 1, 3, 6, and 9 month, the containers of soil were destructively sampled and the dry matter (DM) and WEP of wet soil was determined. The remaining soil sample was air-dried and crushed to pass a 2-mm sieve. Sub-samples were taken, dried at 40°C for 72 h, and analyzed for soil pH and P_m using Morgan's extracting solution. In addition, the 9-month soil samples were analyzed for M3P and M3- Al, Ca, Co, Cu, Fe, K, Mg, Mn, and Zn to determine the effect of amendments on metal and nutrient availability.

Statistical analysis

Data from the four sampling events were analysed as a factorial design with soil type, time and treatment as factors. All interactions were examined. Soil type was a blocking effect in this structure, but its interactions with the randomised factors were of interest. The interactions were interpreted in the first instance by testing simple effects and then making comparisons of means, with adjustment for multiplicity of testing. The GLIMMIX procedure of SAS 9.2 was used to fit the analysis model (SAS 2004). This procedure allowed the addition of heterogeneous variance structures and a number of options for comparing large numbers of means. Residual checks were made to check the assumptions of the analysis method.

Results and Discussion

Slurry and amended slurry analysis

The results of the slurry analysis are shown in Table 2. At the rates used in this study, all of amendments examined reduced the WEP concentration of dairy cattle slurry by approximately 99% compared to the slurry-control ($p < 0.001$). Alum addition reduced slurry pH from approximately 7.2 (slurry-control) to 5.1, PAC reduced pH to 5.7 and FeCl_3 to 5.4 ($p < 0.001$), while lime addition increased slurry pH to 12 ($p < 0.001$). The effectiveness of amendments in reducing P solubility in slurry was in agreement with previous studies (Dao 1999; Lefcourt and Meisinger 2001; Dou et al. 2003; Brennan et al. 2011a). Lefcourt and Meisinger (2001) reported a 97% reduction in WEP of dairy cattle slurry when 2.5% (by weight) of alum was added in a laboratory batch experiment. Dao and Daniel (2002) added alum (810 mg Al L^{-1}) and FeCl_3 (810

mg Fe L⁻¹) (compared to 1,710 mg Al L⁻¹ and 4,560 mg Fe L⁻¹ in the present study) to dairy slurry, and observed that slurry WEP was reduced by 66 and 18%, respectively.

Water extractable phosphorus soil and slurry treatments

There was a significant interaction between soil type, month, and treatment for WEP ($p < 0.001$). The WEP levels from each soil before the start of the incubation period and at each sampling event during the incubation period are shown in Table 3. There was an initial decline in WEP levels for the mineral soils at the 1-month sampling interval (although there was an increase in WEP from Soil E) for the soil-only treatment, followed by an increase at 3 and 6-month sampling intervals, and a trend towards pseudo-steady state by the end of the study. The WEP levels from the soil-only treatment did not vary significantly during the study. As expected, the addition of dairy cattle slurry to soils resulted in observed increases in soil WEP compared to the soil-only treatment for all soils, but these increases in WEP for the slurry treatment were not always statistically significant. There was significantly higher WEP levels following the application of slurry to Soils A, C, and D compared to the soil-only treatment at the 1-month sampling ($p < 0.05$). The WEP levels from the slurry treatment remained significantly higher than the soil-only treatment at the 3-month sampling for Soil C ($p < 0.05$). There were no significant differences between WEP levels for the soil-only and all slurry treatments at the 6 and 9-month sampling events for all four mineral soil types.

The initial reduction in WEP observed in the present study was in agreement with similar incubation studies (Maguire et al. 2001; Griffin et al. 2003; Leytem et al. 2004; Kalbasi and Karthikeyan, 2004; Penn and Bryant, 2006; Dail et al. 2007). Penn and Bryant (2006) examined

the effect of rewetting air-dried sieved soil (2 mm) on WEP and M3P concentration and concluded that decreases in WEP following incubation were likely due to increased microbial activity during initial incubation, resulting in P immobilisation into microbial biomass. The WEP decrease may also have been a result of changes in soil pH, where there was a shift from labile inorganic P to forms that were mineral-associated and less soluble. Changes in pH may have been due to increases in the amount of exchangeable Ca and Mg caused by the breaking down of soil by grinding and sieving. Dail et al. (2007) also found that drying of soil had a significant effect on soil P solubility and concluded that variations due to drying effects should be considered when comparing manures and studies where different drying procedures have been used.

While addition of slurry resulted in an observed increase in WEP for the organic Soil E, the increase was not significant (Table 3). Soil E behaved differently to the mineral soils, and WEP and STP values were typically an order of magnitude greater than for mineral soils. For clarity, the mineral soils and Soil E will be considered separately for the remainder of this paper (Section entitled Organic soils).

Water extractable phosphorus amended slurry treatments

Alum addition decreased WEP of the four mineral soils compared to the slurry-control by between 38 and 97% after 1-month (Table 3) and, on average, by between 52 and 73% for the 4 sampling events. Lime addition decreased WEP by between an average of 59 and 97% after 1-month (50 and 83% for the four sampling events). Poly-aluminium chloride decreased WEP by between 2 and 55% after 1-month. However, PAC increased WEP of Soil B at the 6-month

sampling (Table 3). While this increase was not significant, PAC did not perform as well as alum and lime. Ferric chloride was the least effective at decreasing WEP compared with the other amendments and, in the case of Soil A, B and C, increased WEP compared to the slurry-control. Inherent variability in the consistency of concentrations of TP and WEP within any sample of slurry, combined with different rates in the uptake of slurry P by soil sorption, may complicate interpretation of percentage reductions. These results indicate a difference in effectiveness across soil types and future work must examine the most effective amendments over a much wider range of soil chemical and physical properties.

Water extractable P released (as a percent of total applied P) was used to give a relative comparison of each treatment and soil type after 9-month (Figure 1). The overall treatment trends observed were consistent for the different sampling events of the study and therefore the 9-month values were used as best representing the long-term effect of the amendments on these soils. The treatments with the greatest WEP release compared to the soil-only treatment (in order of decreasing WEP release) were slurry = FeCl_3 > PAC > lime = alum for Soil A and B. For Soils C and D, slurry = FeCl_3 > PAC = alum = lime in terms of WEP release. This shows that alum, lime, and PAC were the most effective amendments at decreasing WEP with slurry-amended soils, while FeCl_3 increased WEP compared to the slurry-control treatment.

In a study that examined the effect of soil P level in a silt loam soil (samples incubated at 25°C), Kalbasi and Karthikeyan (2004) reported that applications of alum and FeCl_3 -amended slurry to soil decreased P solubility, while lime amendments increased WEP. In the present study, alum and FeCl_3 were applied to slurry at lower rates (ratio of 1.5:1 Al:TP and Fe:TP compared to a

ratios of 4:1 and 3.6:1, respectively) than Kalbasi and Karthikeyan (2004). The lower amendment rates used in the present study may somewhat explain the poorer performance of FeCl_3 in reducing WEP levels compared with results from Kalbasi and Karthikeyan (2004), who found FeCl_3 to be the most effective amendment. In addition, the present study used $\text{Ca}(\text{OH})_2$ as the liming product, while Kalbasi and Karthikeyan (2004) used CaO . There were also procedural differences between these two studies: the slurry used by Kalbasi and Karthikeyan (2004) had a much lower DM (2%) than that of the present study (10%), and the slurry application rate used was much lower than in the present study.

In addition to chemical amendment of manure, studies have examined the addition of amendments to high STP soils to reduce P losses (McFarland et al. 2003; Novak and Watts 2005; Brauer et al. 2005). Brauer et al. (2005) incorporated alum ($127 \text{ kg Al ha}^{-1}$) and gypsum at two rates (349 and $1163 \text{ kg Ca ha}^{-1}$) into the upper 10 cm of a high STP soil on an annual basis for 3 yr. Only the high gypsum treatment was observed to reduce WEP and STP values significantly during the study. A limited number of runoff studies have been carried out with chemical amendment of dairy cattle slurry (Elliot et al. 2005; Torbert et al. 2005) and swine slurry (Smith et al. 2001), but little work has focused on the long-term effects of chemical amendments to slurry on the nutrient content of soil.

Soil pH

Within each treatment and soil type, soil pH did not vary significantly with incubation time, with the exception of the FeCl_3 treatment in which soil pH increased with time compared with the other amendments (Table 4). The change in pH may explain the poor performance of FeCl_3 ,

which effectively reduced the WEP of slurry prior to application, but was ineffective when the amended slurry was incubated with soil. There was a strong correlation between soil pH and M3-Al, M3-Ca, M3-Fe, WEP and P_m ($p < 0.001$) for the four mineral soils across all treatments.

The pH of a soil has a significant influence on nutrient availability (Tunney et al. 2010), and changes in pH can alter community composition and activity of microbes in soil (Sylvia et al. 2005). In addition, if the amendments adversely affect microbial activity, the microbes could potentially change the pH by their activity.

Morgan's and Mehlich-3 phosphorus

Figure 2 shows the relationship between P_m and M3P for each treatment at the 9-month sampling event for the mineral soils. There were positive linear relationships between P_m and M3P across all treatments for the mineral soils ($p < 0.01$). The Morgan's P test extracted P from $FeCl_3$ -amended slurry treatment more readily than the M3P test. This was most likely due the difference in pH between Morgan's (4.8) and Mehlich extractants (2.5). These results highlight the importance of selecting the most appropriate extraction methods to assess subsequent soil P availability.

Table 5 shows the temporal variability of P_m over the incubation period for each treatment by soil type combination. There was a significant interaction between soil type, sampling month, and treatment for the P_m of the soil ($p < 0.001$). The P_m of the soil-only treatment did not vary significantly during the course of the experiment. The P_m of the slurry-control remained relatively constant throughout the course of the experiment after the initial increase at 1-month.

The P_m of the slurry-control was significantly different to the soil-only treatment for Soil D (3-month, $p < 0.05$) and Soil C (9-month, $p < 0.15$). These results show that use of chemical amendments did not significantly affect plant availability of soil P. The P_m of the $FeCl_3$ -amended slurry was significantly greater than the slurry-control for Soil A (9-month, $p < 0.05$) and Soil B (6- and 9-month, $p < 0.05$). This was consistent with the results of McFarland et al. (2003) and Kalbasi and Karthikeyan (2004). With the exception of these studies, the authors are not aware of any other studies examining the effect chemical amendment of dairy cattle slurry with alum, $FeCl_3$ and lime on plant availability of P in soils following land application of the slurry.

Mehlich-3 Al, Ca, and Fe analysis

There was a strong correlation between M3-Al and M3-Ca, WEP, pH and P_m ($p < 0.005$); between M3-Ca and pH, WEP, P_m ($p < 0.005$) and between M3-Fe and M3-Al, M3P, and pH ($p < 0.005$) for the four mineral soils (data not shown). When these correlations were examined within treatments, there was no strong correlation between WEP and M3-Al or M3-Fe for any of the slurry amendments. However, the alum treatment tended to increase the levels of M3-Al in these mineral soils, which resulted in largest overall decrease in WEP compared to the lime and PAC treatments. The $FeCl_3$ treatment tended to decrease the M3-Al and M3-Fe levels in these soils, and resulted in an overall increase in WEP levels compared to the slurry-control treatment. There were no observed trends or differences between treatments for M3- Co, Cu, K, Mg, Mn, and Zn. This shows that none of the treatments examined adversely affected the availability of these metals and nutrients to plants.

Relationship between soil WEP and M3P

Figure 3a shows soil M3P at t=0 for pre-incubation soil and at t=9 for all treatments versus WEP for each treatment across the four mineral soils. With the exception of the soil-only treatment, there were significant positive relationships between M3P and WEP for each treatment ($p < 0.05$). Slopes for the alum, PAC, and lime treatments are similar, while the FeCl_3 treatment was steeper, which suggests that for a given rise in M3P, the increase in WEP of the FeCl_3 treatment will be greater compared to the other amendments. Alum, lime and PAC were effective in reducing WEP across the four soil types, which is in agreement with the WEP release results (Table 3).

Relationship between soil WEP and DPS

There was a significant correlation between WEP and DPS for Soil A, Soil B, Soil C and Soil D ($p < 0.05$) (Figure 3b). The WEP-DPS correlation for the soil at t=0 (pre-incubation) and at t=9 month show that as the DPS of the soil increases, the WEP increases. Therefore, amendments which reduce the solubility of P in dairy cattle slurry are effectively increasing the sorption capacity of the soil receiving the amended-slurry. In the present study, the amendments performed differently across different soil types, which indicated that the amendments would be most effective if used in soils with a high DPS. In high DPS soils, chemical amendments will increase the capacity of the soil to store P. There is already an abundance of P sorption sites in soils with low DPS, and apart from the reduction of WEP immediately after slurry application, the long-term benefits would seem to be limited. This work has identified a need to examine further the effect of soil DPS on the effectiveness of chemical amendments in reducing chronic P losses. Soil DPS within a catchment, combined with topography and other P loss risk factors, maybe used to select sites where chemical amendment would be most beneficial.

Organic soils

The organic soil (Soil E) behaved very differently to the mineral soils, as expected (Table 3). None of the amendments effectively reduced the WEP of Soil E (Figure 1). Ann et al. (2000) examined use of chemical amendments to immobilise P in highly organic soils from agricultural land that was being converted into wetlands in the U.S., with the objective of making the wetlands a sink for nutrients. In laboratory-scale experiments, these researchers (Ann et al. 2000) found the order of effectiveness of immobilising P was $\text{FeCl}_3 > \text{alum} > \text{lime}$ when applied at rates of 1-2 g kg⁻¹, 12 g kg⁻¹, and 7-15 g kg⁻¹, respectively, compared to 2.0 g kg⁻¹, 6.6 g kg⁻¹ and 5.6 g kg⁻¹, respectively, in the present study. Ann et al. (2000) concluded that high rates of chemical amendments were needed to reduce P levels because of complexation of P-binding cations (Ca, Fe, and Al) with organic matter. Fox and Kamprath (1971) found that addition of AlCl_2 to an organic soil reduced P solubility significantly. Litaor et al. (2005) showed that drying and re-wetting peat soils resulted in an increased availability of P due to a decrease in P sorption capacity. Litaor et al. (2005) found that rewetting of previously oxidized organic soils dissolved Fe-oxides, resulting in increased P release.

The WEP of Soil E initially increased and remained relatively constant for the duration of the study. The levels of WEP release were as follows: slurry > $\text{FeCl}_3 > \text{lime} > \text{alum} = \text{PAC}$. The differences and trends were not comparable to the mineral soils. In addition, the WEP of the organic soil was an order of magnitude greater than the WEP of the mineral soils, which further complicated comparisons. There was no correlation between P_m and M3P, P_m and WEP, and WEP and M3-Al or M3-Fe for organic soil (data not shown). These results are in agreement with

previous work showing that organic soils have a lower P retention capacity than mineral soils (Iyamuremye and Dick 1996; Daly et al. 2001).

Outlook for use of chemical amendments

Typically, the use of chemically-amended slurry is considered to be most effective within critical source areas (CSAs), where a high STP and a mobilisation vector combine to pose a threat to a nearby receptor. However, the findings of the present study suggest that use of chemically-amended slurry within a CSA could become more refined based on soil suitability criterion. While incubation experiments similar to the type carried out in this study are an accepted means of comparing treatments under similar environmental conditions, it may not be possible to extrapolate the findings of this study to fully represent the effects under field conditions. This study has shown that the effectiveness of amendments is influenced by soil physical and chemical properties. In addition, amendment reduces P solubility in slurry compared to unamended slurry, and offers the opportunity to reduce chronic P losses in addition to their effectiveness in reducing incidental losses.

Future work must examine the long-term effects of amendments at field-scale with a range of soil types. The effect of amendments on P leaching, soil microbiology and structure need to be explored. At present, there is no provision for a licence to land spread any of these amendments in Ireland, except lime to optimize soil pH for production. If chemical amendment were to be used to mitigate P losses, a licensing system would have to be introduced by the Department of Agriculture in Ireland and relevant bodies in other countries. This study indicates that use of chemical amendment as a once-off management practice reduced WEP in soil compared to soil

amended with slurry, but did not decrease STP to levels or have any significant effect on soil physical and chemical properties.

Conclusions

This study found that the WEP of all soils receiving chemically treated dairy cattle slurry was lower than the slurry-control for alum and lime treatments, while FeCl_3 was ineffective at decreasing P solubility. Alum maintained the highest levels of plant available P (M3P) across the four mineral soils with the least risk of P loss to overlying water (i.e., lowest WEP per M3P). This would indicate that alum offers the greatest potential for future research. Further work is needed to examine a much wider range of soil types with a range of DPS with-a-view to identifying soil parameters that could be used to select sites where amendment would be most effective at reducing chronic P losses. The study found that chemical amendments are not effective in organic soils. Future work must examine how amendments affect nutrient balance and pollution swapping at a field-scale under a range of drainage regimes and soil types. While this study has shown that alum is the most effective amendment, further work is required to fully understand the importance of the interactions between the chemistry of amendment and soil, and how such interactions might influence the effectiveness of amendments in reducing chronic P losses. These issues must be addressed before recommendations can be made to farmers or catchment managers.

Acknowledgements

The first author gratefully acknowledges the award of a Walsh Fellowship by Teagasc to support this study. The authors are also grateful for assistance provided by Teagasc and NUI Galway staff and colleagues with special mention to Theresa Cowman, Denis Brennan, Linda Maloney Finn, Tim Sheil, Peter Fahy, Con O'Flynn, Joe Lucid, Stan Lalor, Maja Drapiewska, Carmel O'Connor, and Cecile Labonne.

References

- Andraski, T.W., L.G. Bundy, and K.C. Kilian. 2003. Manure history and long-term tillage effects on soil properties and phosphorus losses in runoff. *Journal of Environmental Quality* 32: 1782–1789.
- Ann, Y., K.R. Reddy, and J.J. Delfino. 1999. Influence of redox potential on phosphorus solubility in chemically amended wetland organic soils. *Ecological Engineering* 14: 169–180.
- Basson, W.D., D.A. Stanton, and R.G., Böhmer. 1968. Automated procedure for the simultaneous determination of phosphorus and nitrogen in plant tissue. *Analyst* 93: 166–172.
- Bond, C.R., R.O. Maguire, and J.L. Havlin. 2006. Change in soluble phosphorus in soils following fertilization is dependent on initial Mehlich-3 phosphorus. *Journal of Environmental Quality* 35:1818-1824.
- Brauer, D., G.E. Aiken, D.H. Pote, S.J. Livingston, L.D. Norton, T.R. Way, and J.H Edwards. 2005. Amendment effects on soil test phosphorus. *Journal of Environmental Quality* 34: 1682-1686.
- Brennan, R.B., O. Fenton, J. Grant, and M.G. Healy. 2011a. Impact of chemical amendment of dairy cattle slurry on phosphorus, suspended sediment and metal loss to runoff from a grassland soil. *Science of the Total Environment* 409: 5111–5118.
- Brennan, R.B., O. Fenton, M. Rodgers, and M.G. Healy. 2011b. Evaluation of chemical amendments to control phosphorus losses from dairy slurry. *Soil Use and Management* 27: 238–246.

Brennan, R.B., M.G. Healy, J. Grant, T.G. Ibrahim, and O. Fenton. 2012. Incidental phosphorus and nitrogen loss from grassland plots receiving chemically amended dairy cattle slurry. *Science of the Total Environment* 441: 132–140.

British Standards, 1990a. British standard methods of test for soils for civil engineering purposes. Determination of particle size distribution. BS 1377:1990:2 BSI, London, United Kingdom.

British Standards, 1990b. Determination by mass-loss on ignition. British standard methods of test for soils for civil engineering purposes. Chemical and electro-chemical tests. BS 1377:1990:3. BSI, London, United Kingdom.

Buda, A.R., P.J.A. Kleinman, M.S. Srinivasan, R.B. Bryant, and G.W. Feyereisen. 2009. Effects of hydrology and field management on phosphorus transport in surface runoff. *Journal of Environmental Quality* 38: 621-627.

Callahan, M.P., P.J.A. Kleinman, A.N. Sharpley, and W.L. Stout, 2002. Assessing the efficacy of alternative phosphorus sorbing soil amendments. *Soil Science* 167: 539-547.

Carpenter, S.R., N.F. Caraco, D.L. Correll, R.W. Howarth, A.N. Sharpley, and V.H. Smith. 1998. Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecological Applications* 8: 559–568.

Coulter, B., and S.T.J. Lalor. 2008. Major and micro nutrient advice for productive agricultural crops. Third edition. Teagasc Johnstown Castle Wexford. 116 pp.

Cummins, T., and E.P. Farrell. 2003. Biogeochemical impacts of clear felling and reforestation on blanket peatland streams I. phosphorus. *Forest Ecology and Management* 180: 545–555.

Dail, H.W., Z. He, M.S. Erich, and C.W. Honeycutt. 2007. 'Effect of Drying on Phosphorus

- Distribution in Poultry Manure', *Communications in Soil Science and Plant Analysis* 38: 13, 1879-1895.
- Daly, K., D. Jeffrey, and H. Tunney. 2001. The effect of soil type on phosphorus sorption capacity and desorption dynamics in Irish grassland soils. *Soil Use and Management* 17: 12–20.
- Dao, T.H. 1999. Co-amendments to modify phosphorus extractability and nitrogen/phosphorus ratio in feedlot manure and composted manure. *Journal of Environmental Quality* 28:1114–1121.
- Dao, T.H., and T.C. Daniel. 2002. Particulate and dissolved phosphorus chemical separation and phosphorus release from treated dairy manure. *Journal of Environmental Quality* 31: 1388-1398.
- Doody, D.G., M. Archbold, R.H. Foy, and R. Flynn. 2012. Approaches to the implementation of the Water Framework Directive: Targeting mitigation measures at critical source areas of diffuse phosphorus in Irish catchments. *Journal of Environmental Management* 93: 225–234.
- Dou, Z., G.Y. Zhang, W.L. Stout, J.D. Toth, and J.D. Ferguson. 2003. Efficacy of alum and coal combustion by-products in stabilizing manure phosphorus. *Journal of Environmental Quality* 32: 1490-1497.
- Elliott, H.A., R.C. Brandt, and G.A. O'Connor. 2005. Runoff phosphorus losses from surface-applied biosolids. *Journal of Environmental Quality* 34: 1632-1639.
- Fox, R.L., and E.J. Kamprath. 1971. Adsorption and leaching of P on acid organic soils and high organic matter sand. *Soil Science Society of America Proceedings* 35: 154-156.
- Griffin, T.S., C.W. Honeycutt, and Z. He. 2003. Changes in soil phosphorus from manure


- application. *Soil Science Society of America Journal* 67: 645–653.
- Iyamuremye, F., and R.P. Dick. 1996. Organic amendments and phosphorus sorption by soils. *Advances in Agronomy* 56: 139–185.
- Kalbasi, M., and K.G. Karthikeyan. 2004. Phosphorus dynamics in soils receiving chemically treated dairy manure. *Journal of Environmental Quality* 33: 2296–2305.
- Kleinman, P., D. Sullivan, A. Wolf, R. Brandt, Z. Dou, H. Elliott, J. Kovar, A. Leytem, R. Maguire, P. Moore, L. Saporito, A. Sharpley, A. Shober, T. Sims, J. Toth, G. Toor, H. Zhang, and T. Zhang. 2007. Selection of a water-extractable phosphorus test for manures and biosolids as an indicator of runoff loss potential. *Journal of Environmental Quality* 36: 1357–1367.
- Kurz, I., H. Tunney, and C.E. Coxon. 2005. The impact of agricultural management practices on nutrient losses to water: data on the effects of soil drainage characteristics. *Water Science and Technology* 51: 73–81.
- Lefcourt, A.M., and J.J. Meisinger. 2001. Effect of adding alum or zeolite to dairy slurry on ammonia volatilization and chemical composition. *Journal of Dairy Science* 84: 1814–1821.
- Leytem, A.B., J.T. Sims, and F.J. Coale. 2004. Determination of phosphorus source coefficients for organic phosphorus sources. *Journal of Environmental Quality* 33: 380–388.
- Litaor, M.I., O. Reichmann, A. Haim, K. Auerswald and M. Shenker. 2005. Sorption characteristics of phosphorus in peat soils of a semi-arid altered wetland. *Soil Science Society of America Journal* 69: 1658-1665.
- McDowell, R.W. and A.N. Sharpley. 2001. Soil phosphorus fractions in solution: influence of fertilizer and manure, filtration and method of determination. *Chemosphere* 45: 737–748.

- McFarland, A.M.S., L.M. Hauck, and A.P. Kruzic. 2003. Phosphorus reductions in runoff and soils from land-applied dairy effluent using chemical amendments: An observation. *Texas Journal of Agriculture and Natural Resources* 16: 47-59.
- Maguire, R.O., and J.T. Sims. 2002. Measuring agronomic and environmental soil phosphorus saturation and predicting phosphorus leaching with Mehlich 3. *Soil Science Society of America Journal* 66: 2033–2039.
- Maguire, R.O., J.T. Sims, S.K. Dentel, F.J. Coale, and J.T. Mah. 2001. Relationships between biosolids treatment process and soil phosphorus availability. *Journal of Environmental Quality* 30:1023-33
- Mehlich, A. 1984. Mehlich 3 soil test extractant: A modification of Mehlich 2 extractant. *Communications in Soil Science and Plant Analysis* 15: 1409–1416.
- Moore, P.A. and D.R. Edwards. 2007. Long-term effects of poultry litter, alum-treated litter, and ammonium nitrate on phosphorus availability in soils. *Journal of Environmental Quality* 36:163–74.
- Morgan, M.F. 1941. Chemical soil diagnosis by the universal soil testing system. Connecticut Agricultural Experiment Station, New Haven, CT. 1941.
- Novak, J.M., and D.W. Watts. 2005. An alum-based water treatment residual can reduce extractable phosphorus concentrations in three phosphorus-enriched coastal plain soils. *Journal of Environmental Quality* 34: 1820-1827.
- O’Flynn, C.J., O. Fenton, P. Wilson, and M.G. Healy. 2012. Impact of pig slurry amendments on phosphorus, suspended sediment and metal losses in laboratory runoff boxes under simulated rainfall. *Journal of Environmental Management* 113: 78–84.

- Penn, C.J., and R.B. Bryant. 2006. Incubation of dried and sieved soils can induce calcium phosphate precipitation/adsorption. *Communications in Soil Science and Plant Analysis* 39: 1437-1449.
- Preedy, N., K. McTiernan, R. Matthews, L. Heathwaite, and P. Haygarth, 2001. Rapid incidental phosphorus transfers from grassland. *Journal of Environmental Quality* 30: 2105–2112.
- Regan, J.T., M. Rodgers, M.G. Healy, L. Kirwan, and O. Fenton. 2010. Determining phosphorus and sediment release rates from five Irish tillage soils. *Journal of Environmental Quality* 39: 1-8.
- SAS. 2004. SAS for Windows. Version 9.1.SAS/STAT® User’s Guide. Cary, NC. SAS Institute Inc.
- Sharpley, A.N. 2003. Soil mixing to decrease surface stratification of phosphorus in manured soils. *Journal of Environmental Quality* 32: 1375–1384.
- Sharpley, A.N., R.W. McDowell, and P.J.A. Kleinman. 2004. Amounts, forms, and solubility of phosphorus in soils receiving manure. *Soil Science Society of America Journal* 68: 2048–2057.
- Sharpley, A.N., T.C. Daniel, J.T. Sims, and D.H. Pote. 1996. Determining environmentally sound soil phosphorus levels. *Journal of Soil and Water Conservation* 51: 160-166.
- Smith, D.R., P.A. Moore, C.L. Griffis, T.C. Daniel, D.R. Edwards, and D.L. Boothe. 2001. Effects of alum and aluminum chloride on phosphorus runoff from swine manure. *Journal of Environmental Quality* 30: 992-998.
- Sylvia, D.M., J.J. Fuhrmann, P. Hartel, and D.A. Zuberer. 2005. Principles and applications of soil microbiology. Pearson Prentice Hall, New Jersey.


- Torbert, H.A., K.W. King, and R.D., Harmel. 2005. Impact of soil amendments on reducing phosphorus losses from runoff in sod. *Journal of Environmental Quality* 34: 1415-1421.
- Tunney, H. 2000. Phosphorus needs of grassland soils and loss to water. In: Steenvoorden, J., Claesses, F., Willems, J. (eds.). International conference on agricultural effects on ground and surface waters. International association of hydrological sciences, Wageningen, Netherlands. pp. 63-69.
- Tunney, H., L., Kirwan, W. Fu, N. Culleton, and A.D. Black. 2010. Long-term phosphorus grassland experiment for beef production - impacts on soil phosphorus levels and live weight gains. *Soil Use and Management* 26: 237-244.
- Wall, D.P., P. Jordan, A.R. Melland, P-E. Mellander, S. Mechan, and G. Shortle. 2013. Forecasting the decline of excess soil phosphorus in agricultural catchments. *Soil Use and Management* 29: 147-154.

Figure 1. Water extractable phosphorus release (WEP) expressed as a percentage of total phosphorus applied for nine month compared to soil-only treatment (a) the four mineral soils (b) peat soil (PAC, poly-aluminum chloride)



PAC, poly-aluminum chloride

Figure 2. Morgan's P (P_m) against Mehlich-3 extractable phosphorus (M3P) for the four mineral soils by treatment (nine-month data)



PAC, poly-aluminum chloride

Figure 3. Soil water extractable phosphorus (WEP) against (a) Mehlich-3 extractable phosphorus (M3P) and (b) degree of phosphorus saturation (DPS) for the four mineral soils (nine month data)




Table 1**Soil location, texture and physical and chemical properties**

| Site | | Soil A | Soil B | Soil C | Soil D | Soil E |
|---|--------------------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| Location | | Cork | Wexford | Wexford | Galway | Sligo |
| Coordinates | | 52°07' N, 8°16' W | 52°17' N, 6°31' W | 52°17' N, 6°31' W | 53°21' N, 8°34' W | 54°04' N, 8°52' W |
| Soil texture | | Sandy loam | Clay loam | Loam | Silty loam | - |
| Sand content | % | 56.2 (1.1) | 51.8 (4.2) | 37.8 (1.1) | 15.0 (1.4) | - |
| Silt content | % | 25.8 (1.3) | 28.1 (4.9) | 31.1 (1.0) | 72.0 (1.1) | - |
| Clay content | % | 18.0 (2.4) | 20.1 (2.2) | 31.1 (2.1) | 13.0 (1.2) | - |
| Organic matter | % | 7.9 (0.6) | 7.8 (0.3) | 6.7 (0.5) | 13.3 (0.2) | 77.4 (0.4) |
| Soil pH | | 6.1 (0.9) | 5.7 (0.1) | 6.50 (0.02) | 5.10 (0.04) | 5.6 (0.1) |
| P Index ^a | Index | 3 | 3 | 1 | 3 | 4 |
| Water extractable phosphorus (WEP) | mg kg ⁻¹ | 7.2 (1.2) | 6.2 (2.1) | 2.7 (0.9) | 3.2 (1.5) | 42.5 (4.5) |
| Morgan's extractable phosphorus (P _m) | mg L ⁻¹ | 5.8 (0.3) | 5.7 (0.1) | 2.6 (0.2) | 5.1 (0.4) | 24.6 (0.2) |
| Degree of phosphorus saturation (DPS) | % | 3.94 (0.39) | 5.62 (0.09) | 1.52 (0.07) | 3.34 (0.13) | 17.3 (0.2) |
| Mehlich-3 extractable phosphorus (M3P) | mg kg ⁻¹ | 47.3 (2.4) | 68.7 (1.2) | 16.2 (0.9) | 34.7 (1.1) | 121.0 (5.4) |
| Mehlich-3 extractable aluminum (M3-Al) | mg kg ⁻¹ | 765 (60) | 871 (9) | 969 (4) | 620 (17) | 304 (24) |
| Mehlich-3 extractable copper (M3-Ca) | mg kg ⁻¹ | 1780 (57) | 1320 (37) | 1340 (9) | 258 (19) | 657 (117) |
| Mehlich-3 extractable cobalt (M3-Co) | mg kg ⁻¹ | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Mehlich-3 extractable copper (M3-Cu) | mg kg ⁻¹ | 2.7 (0.7) | 3.1 (0.2) | 3.4 (0.0) | 4.5 (0.1) | 1.8 (0.1) |
| Mehlich-3 extractable iron (M3-Fe) | mg kg ⁻¹ | 516 (2) | 447 (10) | 213 (2) | 479 (6) | 444 (6) |
| Mehlich-3 extractable potassium (M3-K) | mg kg ⁻¹ | 168 (4) | 133 (4) | 270 (5) | 153 (7) | 336 (3) |
| Mehlich-3 extractable magnesium (M3-Mg) | mg kg ⁻¹ | 199 (5) | 203 (6) | 305 (5) | 122 (3) | 1110 (7) |
| Mehlich-3 extractable molybdenum (M3-Mn) | mg kg ⁻¹ | 93 (5) | 102 (1) | 127 (13) | 75.3 (0.6) | 14.9 (1.1) |
| Mehlich-3 extractable zinc (M3-Zn) | mg kg ⁻¹ | 2.3 (0.1) | 4.8 (0.0) | 3.5 (0.9) | 3.7 (0.1) | 17.1 (3.6) |
| Container capacity (CC) | kg water kg soil ⁻¹ | 675 (32) | 634 (12) | 539 (8) | 825 (43) | 2110 (59) |
| Soil bulk density | kg m ⁻³ | 1.29 | 1.15 | 1.08 | 0.93 | 0.27 |

() standard deviations; ^aP Index: soils in Ireland ranked based on risk of P loss (Index 1 being low risk and 4 being the highest)

Table 2
Slurry and amended slurry properties (standard deviation in brackets)

| Treatment | DM % | pH | TN mg L ⁻¹ | TP mg L ⁻¹ | WEP mg kg ⁻¹ | TK mg L ⁻¹ |
|-------------------|--------------|------------|--------------------------|--------------------------|----------------------------|--------------------------|
| Slurry-control | 10.45 (0.78) | 7.2 (0.3) | 4860 (425) | 1140 (93) | 2.86 (0.42) | 3110 (254) |
| Alum | 10.1 (0.4) | 5.1 (0.4) | 4660 (201) | 1120 (19) | 0.003 (0.001) | 2840 (167) |
| Lime | 13.8 (0.2) | 12.0 (0.3) | 3590 (459) | 944 (79) | 0.025 (0.001) | 2620 (430) |
| PAC | 9.9 (0.4) | 5.7 (0.4) | 5320 (379) | 1280 (154) | 0.003 (0.001) | 3060 (617) |
| FeCl ₃ | 11.4 (0.7) | 5.4 (0.1) | 5020 (283) | 1180 (84) | 0.020 (0.005) | 3100 (153) |

() standard deviation; DM, dry matter; TN, total nitrogen; TP, total phosphorus; WEP, water extractable phosphorus; TK, total potassium; PAC, poly-aluminium chloride.

Table 3

Water extractable phosphorus concentration (WEP) and percentage decrease for each treatment at each sampling event

| Soil | Month | Slurry-control | | | | Lime | | PAC | | FeCl ₃ | |
|------|-------|----------------------------------|--------------------------|-----------------------------|----|-------------------------|-----|---------------------------|-----|-------------------------|-----|
| | | Soil-only mg kg ⁻¹ | mg kg ⁻¹ | Alum mg kg ⁻¹ | % | mg kg ⁻¹ | % | mg kg ⁻¹ | % | mg kg ⁻¹ | % |
| A | 0 | 7.2 (0.9) | | | | | | | | | |
| | 1 | 2.3 (0.3) | 4.1 (1.5) | 2.4 (0.1) | 38 | 1.6 (1.0) | 59 | 4 (2) | 2 | 5.0 (3.1) | -26 |
| | 3 | 4.6 (1.0) | 19.9 (3.4)* ^a | 4.8 (0.5)* ^b | 75 | 7.5 (1.5) | 61 | 8.9 (1.1) | 55 | 2.2 (1.6)* ^b | 88 |
| | 6 | 3.4 (0.3) | 8.4 (0.3) | 4.0 (0.2) | 51 | 4.1 (0.3) | 51 | 6.2 (0.7) | 25 | 8.2 (4.2) | 2 |
| | 9 | 3.3 (1.1) | 6.1 (1.0) | 3.3 (0.2) | 44 | 4.0 (0.5) | 33 | 5.2 (0.4) | 13 | 10.2 (0.3) | -71 |
| B | 0 | 6.2 (1.3) | | | | | | | | | |
| | 1 | 4.7 (0.7) | 9.1 (2.7) | 1.0 (1.4)* ^b | 88 | 0.5 (0.3)* ^b | 93 | 2.8 (1.6)* ^b | 68 | 4.5 (1.9) | 50 |
| | 3 | 5.8 (1.4) | 13.0 (2.4) | 8.4 (2.0) | 38 | 10.2 (2.6) | 24 | 9.8 (1.3) | 27 | 15.2 (2.4) | -12 |
| | 6 | 6.3 (4.1) | 8.7 (0.1) | 4.8 (0.4) | 43 | 4.9 (0.3) | 42 | 11.0 (0.2) | -28 | 1.3 (0.0) | 84 |
| | 9 | 3.0 (0.3) | 7.9 (1.3) | 3.4 (0.9) | 56 | 4.9 (1.1) | 38 | 5.8 (0.4) | 26 | 13.4 (0.2) | -70 |
| C | 0 | 2.6 (0.7) | | | | | | | | | |
| | 1 | 0.1 (0.0) | 7.3 (1.2)* ^a | 0.10 (0.04)* ^b | 97 | 0.1 (0.0)* ^b | 97 | 0.10 (0.01)* ^b | 97 | 1.4 (2.2)* ^b | 80 |
| | 3 | 0.6 (0.1) | 5.4 (1.3)* ^a | 3.1 (0.6) | 42 | 0.6 (0.2)* ^b | 88 | 2.6 (1.9) | 52 | 10.3 (0.9) | -92 |
| | 6 | 5.5 (0.5) | 8.8 (1.8) | 1.4 (0.2) | 83 | 2.3 (0.4) | 73 | 5.7 (1.4) | 35 | 2.5 (0.2) | 71 |
| | 9 | 1.3 (0.1) | 5.7 (0.3)* ^a | 2.2 (0.6) | 61 | 1.6 (0.5) | 72 | 1.7 (0.5) | 69 | 5.9 (0.3) | -5 |
| D | 0 | 3.2 (1.3) | | | | | | | | | |
| | 1 | 0.2 (0.1) | 6.0 (2.2) * ^a | 0.3 (0.2)* ^b | 94 | 0.2 (0.1)* ^b | 96 | 0.5 (0.4)* ^b | 91 | 2.8 (1.9) | 53 |
| | 3 | 4.8 (0.3) | 10.0 (0.7) | 0.60 (0.03)* ^b | 94 | 2.3 (3.1)* ^b | 77 | 0.50 (0.02) | 94 | 8.2 (2.8) | 21 |
| | 6 | 2.9 (0.2) | 10.4 (0.9) | 5.2 (3.2) | 50 | 4.3 (1.1) | 58 | 7.5 (1.0) | 27 | 7.4 (1.4) | 28 |
| | 9 | 2.3 (0.2) | 6.2 (0.6) | 2.8 (1.2) | 53 | 3.4 (0.2) | 45 | 3.1 (0.7) | 50 | 5.9 (0.5) | 4 |
| E | 0 | 43.0 (4.4) | | | | | | | | | |
| | 1 | 46.9 (8.5) | 63.8 (9.0) | 51 (12) | 19 | 56.9 (7.3) | 10 | 44.8 (1.12) | 29 | 67.5 (3.5) | -7 |
| | 3 | 61.8 (8.5) | 104 (7) | 92 (16) | 10 | 74.5 (12.6) | 27 | 69.0 (13.9) | 33 | 90.4 (17.4) | 12 |
| | 6 | 63.1 (4.2) | 88.6 (8.9) | 78.6 (9.4) | 10 | 97.1 (13) | -11 | 64.3 (1.7) | 26 | 109 (13) | -24 |
| | 9 | 48.0 (3.4) | 64.9 (0.3) | 53 (8) | 17 | 64.2 (3.3) | 0 | 55.0 (12.6) | 14 | 85.1 (6.1) | -32 |

() standard deviation; *indicate statistical significance of 0.05; ^acompared with soil-only treatment; ^bcompared to slurry-control; PAC, poly-aluminium chloride

1 **Table 4** Soil pH for each treatment at each sampling event

| Soil | Month | Soil-only mg kg ⁻¹ | Slurry-control mg kg ⁻¹ | Alum mg kg ⁻¹ | Lime mg kg ⁻¹ | PAC mg kg ⁻¹ | FeCl ₃ mg kg ⁻¹ |
|------|-------|----------------------------------|---------------------------------------|-----------------------------|-----------------------------|----------------------------|--|
| A | 0 | 6.1 (0.9) | | | | | |
| | 1 | 5.40 (0.02) | 5.5 (0.1) | 5.10 (0.01) | 5.8 (0.7) | 5.50 (0.01) | 6.3 (0.8) |
| | 3 | 5.5 (0.0) | 6.2 (0.7) | 5.6 (0.1) | 6.4 (0.8) | 5.7 (0.4) | 6.2 (1.1) |
| | 6 | 5.20 (0.03) | 5.10 (0.02) | 4.70 (0.02) | 5.00 (0.01) | 4.98 (0.01) | 6.1 (1.1) |
| | 9 | 5.30 (0.06) | 5.20 (0.05) | 4.70 (0.06) | 5.00 (0.04) | 5.00 (0.02) | 6.9 (0.1)* ^{a,b} |
| B | 0 | 5.70 (0.08) | | | | | |
| | 1 | 5.00 (0.01) | 5.20 (0.03) | 5.00 (0.02) | 5.40 (0.02) | 5.6 (0.1) | 6.2 (0.2) |
| | 3 | 5.9 (0.7) | 6.0 (1.2) | 5.9 (0.7) | 5.8 (0.9) | 5.5 (0.3) | 5.5 (0.2) |
| | 6 | 5.4 (0.1) | 4.8 (0.1) | 4.50 (0.03) | 4.80 (0.02) | 5.3 (0.8) | 6.3 (0.0) |
| | 9 | 5.0 (0.1) | 4.80 (0.02) | 4.50 (0.06) | 4.80 (0.02) | 4.80 (0.02) | 6.6 (0.2) |
| C | 0 | 6.50 (0.02) | | | | | |
| | 1 | 6.30 (0.03) | 6.20 (0.04) | 5.60 (0.01) | 6.1 (1.1) | 6.30 (0.04) | 7.3 (0.1) |
| | 3 | 6.1 (0.4) | 5.7 (0.2) | 5.6 (0.5) | 6.8 (1.1) | 5.8 (0.5) | 6.1 (0.9) |
| | 6 | 5.5 (0.5) | 5.9 (0.4) | 5.6 (0.1) | 5.7 (0.1) | 6.3 (0.9) | 6.6 (1.4) |
| | 9 | 6.30 (0.03) | 6.10 (0.03) | 5.40 (0.03) | 5.60 (0.03) | 5.70 (0.03) | 7.3 (0.1) |
| D | 0 | 5.10 (0.04) | | | | | |
| | 1 | 5.30 (0.02) | 5.60 (0.05) | 5.50 (0.06) | 6.10 (0.03) | 6.3 (0.0) | 6.7 (0.1)* ^a |
| | 3 | 5.60 (0.05) | 6.5 (1.1) | 5.50 (0.01) | 6.1 (0.7) | 5.9 (0.3) | 5.9 (0.6) |
| | 6 | 5.1 (0.1) | 5.0 (0.3) | 5.0 (0.2) | 5.2 (0.1) | 5.5 (0.5) | 6.0 (0.7) |
| | 9 | 5.00 (0.05) | 5.0 (0.1) | 4.8 (0.1) | 5.0 (0.1) | 5.1 (0.1) | 6.4 (0.1) |
| E | 0 | 5.60 (0.05) | | | | | |
| | 1 | 5.8 (0.2) | 5.70 (0.06) | 5.60 (0.02) | 5.70 (0.01) | 6.00 (0.04) | 5.50 (0.02) |
| | 3 | 5.3 (0.2) | 5.3 (0.1) | 5.3 (0.2) | 5.4 (0.2) | 5.4 (0.3) | 5.70 (0.01) |
| | 6 | 5.20 (0.04) | 5.1 (0.1) | 5.09 (0.05) | 5.3 (0.3) | 5.5 (0.2) | 5.20 (0.01) |
| | 9 | 5.3 (0.1) | 5.10 (0.01) | 5.1 (0.1) | 5.10 (0.05) | 5.7 (0.2) | 5.20 (0.04) |

2 () standard deviation; *indicate statistical significance of 0.05; ^acompared with soil-only treatment; ^bcompared to slurry-control; PAC, poly-aluminium chloride.

3 **Table 5** Morgan's extractable phosphorus (P_m) for each treatment at each sampling event

| Soil | Month | Soil-only mg kg ⁻¹ | Slurry-control mg kg ⁻¹ | Alum mg kg ⁻¹ | Lime mg kg ⁻¹ | PAC mg kg ⁻¹ | FeCl ₃ mg kg ⁻¹ |
|------|-------|----------------------------------|---------------------------------------|-----------------------------|-----------------------------|----------------------------|--|
| A | 0 | 6 (1) | | | | | |
| | 1 | 5.9 (0.2) | 11.9 (0.9) | 6.1 (0.6) | 8.2 (2.8) | 7.7 (0.6) | 10.9 (4.1) |
| | 3 | 2.6 (0.1) | 10.0 (0.6) | 4.0 (0.9) | 5.7 (2.0) | 5.8 (3.0) | 10.1 (2.6)* ^a |
| | 6 | 6.2 (0.3) | 12.2 (0.5) | 6.8 (0.4) | 6.8 (0.2) | 8.6 (0.3) | 21.4 (4.8) |
| | 9 | 6.2 (0.1) | 12.2 (0.1) | 8.3 (0.1) | 7.2 (0.4) | 9.2 (0.8) | 26.8 (2.5)* ^{a,b} |
| B | 0 | 5.7 (0.1) | | | | | |
| | 1 | 6.3 (0.2) | 10.4 (1.5) | 6.8 (0.2) | 7.1 (0.1) | 7.6 (0.1) | 12.1 (2.2) |
| | 3 | 7.0 (0.4) | 15.0 (3.8) | 7.9 (1.1) | 8.3 (1.0) | 8.0 (0.6) | 8.9 (3.2) |
| | 6 | 4.8 (0.1) | 11.1 (0.0) | 7.6 (0.4) | 7.8 (0.1) | 8.3 (0.3) | 23.2 (3.3)* ^{a,b} |
| | 9 | 6.0 (0.1) | 11.9 (0.4) | 9.0 (0.3) | 8.5 (0.4) | 9.2 (0.3) | 26.8 (5.6)* ^{a,b} |
| C | 0 | 2.6 (0.2) | | | | | |
| | 1 | 2.2 (0.2) | 6.6 (1.1) | 2.7 (0.4) | 2.5 (0.3) | 3.5 (1.0) | 8.6 (4)* ^a |
| | 3 | 2.4 (0.1) | 6.1 (0.8) | 7.2 (4.9) | 3.9 (1.5) | 5.7 (1.5) | 14.5 (1.9)* ^a |
| | 6 | 2.5 (0.5) | 6.8 (0.6) | 2.7 (0.2) | 2.8 (0.6) | 6.8 (5.6) | 9.1 (4.2) |
| | 9 | 2.2 (0.2) | 6.7 (0.4) | 3.6 (0.1) | 2.9 (0.4) | 4.1 (0.4) | 14.4 (2.3)* ^{a,b} |
| D | 0 | 5.1 (0.4) | | | | | |
| | 1 | 4.7 (0.4) | 9.0 (0.9) | 5.4 (0.4) | 6.0 (0.3) | 7.8 (1.0) | 12.7 (2.9) |
| | 3 | 5.0 (0.1) | 9.9 (4.4)* ^a | 5.6 (0.0) | 6.7 (2.7) | 6.7 (0.8) | 11.7 (4.0) |
| | 6 | 4.6 (0.7) | 7.5 (1.8) | 5.1 (0.2) | 6.1 (0.0) | 9.6 (5.5) | 13.3 (1.0) |
| | 9 | 5.3 (0.1) | 8.1 (0.5) | 5.3 (1.0) | 5.3 (0.6) | 6.2 (0.4) | 15.9 (1.4)* ^a |
| E | 0 | 24.6 (0.2) | | | | | |
| | 1 | 34.6 (4.1) | 43.8 (1.4) | 38.1 (1.2) | 44.1 (1.0) | 48.9 (3.6) | 49.5 (6.4) |
| | 3 | 38.3 (1.2) | 35.3 (1.3) | 40.1 (7.8) | 37.8 (0.9) | 45.3 (1.9) | 39 (7.1) |
| | 6 | 33.0 (3.4) | 37.0 (2.1) | 29.9 (1.5) | 28.5 (1.3) | 31.9 (1.1) | 35.3 (0.6) |
| | 9 | 28.8 (5.4) | 40.1 (2.0) | 35.9 (1.2) | 39.6 (3.5) | 40.1 (7.3) | 42.3 (3.9) |

() standard deviation; *indicate statistical significance of 0.05; ^acompared with soil-only treatment; ^bcompared to slurry-control; PAC, poly-aluminium chloride.

4
5
6