



Provided by the author(s) and University of Galway in accordance with publisher policies. Please cite the published version when available.

Title	Occupational therapy-led interventions for people with anxiety disorders - impact on functioning and mental health symptoms
Author(s)	Fox, Jackie; Erlandsson, Lena-Karin; Shiel, Agnes
Publication Date	2018-06-11
Publication Information	Fox, Jackie, Erlandsson, Lena-Karin , & Shiel, Agnes. (2018). Occupational therapy-led interventions for people with anxiety disorders - impact on functioning and mental health symptoms Paper presented at the RCOT2018 Annual Conference and Exhibition, Belfast, 11-13 June.
Publisher	Royal College of Occupational Therapists
Link to publisher's version	https://rcotannualconference.org.uk/wp-content/uploads/2018/06/ABSTRACTS-2018-ONLINE.pdf
Item record	http://hdl.handle.net/10379/7488

Downloaded 2024-05-05T03:59:21Z

Some rights reserved. For more information, please see the item record link above.





Occupational therapy-led interventions for people with anxiety disorders

Impact on functioning and mental health symptoms

Jackie Fox^a, Lena-Karin Erlandsson^b & Agnes Shiel^a

^a *School of Health Sciences, National University of Ireland, Galway.*

^b *Department of Health Science, Lund University, Sweden*

Anxiety is prevalent and is associated with loss of functioning in occupations like work and household management (Kessler et al., 2009). But the effectiveness of occupational therapy interventions to improve functioning for this population was under-researched. This systematic review aimed to determine the effectiveness of occupational therapy interventions to improve functioning and mental health outcomes for individuals with anxiety disorders.


The systematic review included studies involving (a) individuals with anxiety and stress-related disorders over 18 years, (b) outcomes relating to functioning and/or mental health and (c) interventions designed/led/facilitated by an occupational therapist. Eleven databases were searched for literature published from 1994 – December 2017. Data extraction was performed by the first author and independently reviewed by the other authors. The studies were critically appraised for methodological quality (Law & MacDermid, 2014).

The review found 19 papers describing 13 individual studies. Because the nature of the interventions varied, narrative analysis was used rather than meta-analysis. Interventions were clustered into discrete categories depending on the underpinning theory of the intervention, namely (i) cognitive-behavioural theory, (ii) lifestyle modification, (iii) skill-building, (iv) the ValMO model (Persson, Erlandsson, Eklund, & Iwarsson, 2001) and (v) neurological and sensory. Interventions using lifestyle modification (Lambert et al., 2007), the ValMO model (Eklund & Erlandsson, 2011) and skill-building (Helfrich, Peters, & Chan, 2011) showed the potential for effectiveness in improving both anxiety symptoms and daily functioning.

Many reviewed studies were small and uncontrolled, indicating an urgency for high quality research in this area. Occupational therapy interventions have the potential to improve functioning and mental health outcomes for people with anxiety, but interventions require replication and larger-scale research to demonstrate the added value of occupational therapy (Ashby, Gray, Ryan, & James, 2015).

References (used in abstract)

- Ashby, S., Gray, M., Ryan, S., & James, C. (2015). Maintaining occupation-based practice in Australian mental health practice: A critical stance. *British Journal of Occupational Therapy*, 78(7), 431-439.
- Eklund, M., & Erlandsson, L.-K. (2011). Return to work outcomes of the Redesigning Daily Occupations (ReDO) program for women with stress-related disorders—A comparative study. *Women & Health*, 51(7), 676-692.
- Helfrich, C. A., Peters, C. Y., & Chan, D. V. (2011). Trauma symptoms of individuals with mental illness at risk for homelessness participating in a life skills intervention. *Occupational Therapy International*, 18(3), 115-123. doi: <http://dx.doi.org/10.1002/oti.308>
- Kessler, R. C., Aguilar-Gaxiola, S., Alonso, J., Chatterji, S., Lee, S., & Ustun, T. B. (2009). The WHO World Mental Health (WMH) Surveys. *Psychiatrie (Stuttg)*, 6(1), 5-9.
- Law, M., & MacDermid, J. (Eds.). (2014). *Evidence-Based Rehabilitation: A Guide to Practice* (3rd ed.). Thorofare, NJ: SLACK Incorporated.
- Persson, D., Erlandsson, L.-K., Eklund, M., & Iwarsson, S. (2001). Value dimensions, meaning, and complexity in human occupation—a tentative structure for analysis. *Scandinavian Journal of Occupational Therapy*, 8(1), 7-18.



Papers included in the final review, clustered by theoretical underpinning:

Interventions using cognitive-behavioural theory

- Kitchiner, N. J., Edwards, D., Wood, S., Sainsbury, S., Hewin, P., Burnard, P., & Bisson, J. I. (2009). A randomized controlled trial comparing an adult education class using cognitive behavioural therapy "stress control", anxiety management group treatment and a waiting list for anxiety disorders. *Journal of Mental Health, 18*(4), 307-315.
- Kohn, M., Hitch, D., & Stagnitti, K. (2012). Better Access to Mental Health program: influence of mental health occupational therapy. *Australian Occupational Therapy Journal, 59*(6), 437-444. doi: <http://dx.doi.org/10.1111/1440-1630.12005>
- Prior, S. (1998a). Anxiety Management: Results of a Follow-Up Study. *British Journal of Occupational Therapy, 61*(6), 284-285.
- Prior, S. (1998b). Determining the Effectiveness of a Short-Term Anxiety Management Course. *British Journal of Occupational Therapy, 61*(5), 207-213.

Interventions using lifestyle modification theory

- Lambert, R. A., Harvey, I., & Poland, F. (2007). A pragmatic, unblinded randomised controlled trial comparing an occupational therapy-led lifestyle approach and routine GP care for panic disorder treatment in primary care. *Journal of Affective Disorders, 99*(1-3), 63-71.
- Lambert, R., Caan, W., & McVicar, A. (2008). Influences of lifestyle and general practice (GP) care on the symptom profile of people with panic disorder. *Journal of Public Mental Health, 7*(2), 18-24. doi: 10.1108/17465729200800011

Interventions using skill-building theory (specifically for those with PTSD or trauma)

- Classen, S., Cormack, N. L., Winter, S. M., Monahan, M., Yarney, A., Lutz, A. L., & Platek, K. (2014). Efficacy of an occupational therapy driving intervention for returning combat veterans. *OTJR Occupation, Participation and Health, 34*(4), 176-182.
- Classen, S., Monahan, M., Canonizado, M., & Winter, S. (2014). Utility of an Occupational Therapy Driving Intervention for a Combat Veteran. *American Journal of Occupational Therapy, 68*(4), 405-411. doi: 10.5014/ajot.2014.010041
- Helfrich, C. A., Peters, C. Y., & Chan, D. V. (2011). Trauma symptoms of individuals with mental illness at risk for homelessness participating in a life skills intervention. *Occupational Therapy International, 18*(3), 115-123. doi: <http://dx.doi.org/10.1002/oti.308>
- Phillips, M. E., Bruehl, S., & Harden, R. (1997). Work-related post-traumatic stress disorder: Use of exposure therapy in work-simulation activities. *American Journal of Occupational Therapy, 51*(8), 696-700. doi: <http://dx.doi.org/10.5014/ajot.51.8.696>
- Rogers, C. M., Mallinson, T., & Peppers, D. (2014). High-intensity sports for posttraumatic stress disorder and depression: Feasibility study of ocean therapy with veterans of operation enduring freedom and operation Iraqi freedom. *American Journal of Occupational Therapy, 68*(4), 395-404.

Interventions using occupational science theory (the ValMO model)

- Eklund, M. (2013). Anxiety, depression, and stress among women in work rehabilitation for stress-related disorders. *International Journal of Mental Health, 42*(4), 34-47. doi: 10.2753/imh0020-7411420402
- Eklund, M., & Erlandsson, L.-K. (2011). Return to work outcomes of the Redesigning Daily Occupations (ReDO) program for women with stress-related disorders—A comparative study. *Women & Health, 51*(7), 676-692.
- Eklund, M., & Erlandsson, L.-K. (2013). Quality of life and client satisfaction as outcomes of the Redesigning Daily Occupations (ReDO) programme for women with stress-related disorders: A comparative study. *Work, 46*(1), 51-58.

Eklund, M., & Erlandsson, L.-K. (2014). Women's perceptions of everyday occupations: outcomes of the Redesigning Daily Occupations (ReDO) programme. *Scandinavian Journal of Occupational Therapy*, 21(5), 359-367. doi: 10.3109/11038128.2014.922611

Eklund, M., Wästberg, B. A., & Erlandsson, L.-K. (2013). Work outcomes and their predictors in the Redesigning Daily Occupations (Re DO) rehabilitation programme for women with stress-related disorders. *Australian Occupational Therapy Journal*, 60(2), 85-92. doi: 10.1111/1440-1630.12019

Interventions using neurological/sensory theories

Bracciano, A. G., Chang, W. P., Kokesh, S., Martinez, A., Meier, M., & Moore, K. (2012). Cranial Electrotherapy Stimulation in the Treatment of Posttraumatic Stress Disorder: A Pilot Study of Two Military Veterans. *Journal of Neurotherapy*, 16(1), 60-69.

Champagne, T. (2011). The influence of posttraumatic stress disorder, depression, and sensory processing patterns on occupational engagement: a case study. *Work*, 38(1), 67-75 69p. doi: 10.3233/wor-2011-1105

Moore, K. M., & Henry, A. D. (2002). Treatment of adult psychiatric patients using the Wilbarger protocol. *Occupational Therapy in Mental Health*, 18(1), 43-63.

Citation: Fox, J., Erlandsson, L.K. & Shiel, A. (2018). *Occupational therapy-led interventions for people with anxiety disorders: Impact on functioning and mental health symptoms*. Paper presented at the Royal College of Occupational Therapists Annual Conference, Belfast, UK.

Further information on upcoming publication of this work can be found at:

https://www.researchgate.net/profile/Jackie_Fox?ev=hdr_xprf&_sg=P5n_PTLJ1j9hni2ZwX84eJ3HFaiWBecPtMMAT052Um01UBhfxlep78x7_2ZRbTuQvbpjSkKOoAqXoNmPw_n316yZ

or from the author at jackie.fox@nuigalway.ie